

## Mathematical model limits malaria outbreaks

## January 24 2017

One of the most common infectious diseases in the world, malaria causes public health problems and depresses the economy of infected areas. When untreated or treated improperly, the disease can result in fatalities. Despite impressive control measures and increased prevention techniques, which have reduced the global malaria mortality rate by 29% over the last six years, 3.3 billion people throughout 97 countries and territories still face a risk of infection. According to the World Health Organization, there were 212 million cases of malaria in 2015; approximately 429,000 resulted in death. Sub-Saharan Africa continues to exhibit a disproportionately high number of outbreaks and fatalities.

Mathematical models can effectively predict and track <u>malaria</u> transmission trends, ultimately quantifying the efficiency of various treatment and eradication strategies in high-risk regions. In a paper publishing in the *SIAM Journal on Applied Mathematics* on January 24th, Xiunan Wang and Xiao-Qiang Zhao explain a malaria transmission model that considers three distinct factors: climate, the extrinsic incubation period (EIP), and the vector-bias effect. Using data from Maputo Province, Mozambique to simulate transmission trends, the authors ultimately present a possible way to limit the disease's transmission.

Female Anopheles mosquitoes are responsible for the transmission of malaria, which is caused by the one-celled Plasmodium parasite. A variety of environmental factors—including temperature, rainfall, humidity, and wind patterns—significantly impact the maturity,



reproduction, and longevity of mosquitoes. The mosquito life cycle, in turn, directly affects the parasite's survival. "Climate factors have great impact on mosquito life cycle and parasite development," Wang said. "It becomes particularly important to consider climate impact on malaria transmission in light of global climate change." For example, an increase in temperature lessens the number of days necessary for breeding and quickens formation of spores in the parasites. Both of these occurrences can increase transmission.

The EIP is the length of time a parasite needs to fully develop in a mosquito and migrate to its salivary glands in preparation for transmission. Female mosquitoes live anywhere from three to 100 days, and a typical EIP ranges from 10 to 30 days. "Only those mosquitoes that live long enough to survive the EIP can transmit malaria," Wang said. They are infectious for the rest of their lives. Additionally, previous research confirmed that malaria-infected humans attract more mosquitoes due to chemical substances emitted by the parasite. Therefore, it seems that mosquitoes prefer infected hosts rather than susceptible ones—a skewed probability of infection known as "vector bias."

Wang and Zhao adopt prior malaria transmission models in their own vector-bias model, which utilizes the basic reproduction ratio R0 and incorporates the aforementioned three factors. "The basic reproduction ratio serves as a threshold parameter in determining the global stability of either disease-free or endemic periodic solutions for this period and time-delayed system," Wang said. She and Zhao treat all parameters related to humans as constants but assume that mosquito-related parameters are periodic functions, thus incorporating seasonality into the model. Two additional parameters quantify the vector-bias effect, the probability of a mosquito biting an infected or susceptible human. And a constant time delay represents the EIP. The authors apply their model to published data from Maputo Province and simulate transmission trends



in the area; the simulated curve matches the real-data curve for transmission. "The numerical simulations for monthly new malaria cases are well consistent with the real data from Maputo Province," Wang said. "This suggests that such a model may give a more accurate prediction of the disease <u>transmission</u>."

Based on their findings of the model's global dynamics, Wang and Zhao discover that malaria will continue to exhibit seasonal fluctuation in Maputo Province in the coming years. More importantly, they show that a shorter EIP directly corresponds with increased <u>malaria transmission</u>. Thus, extending the EIP of mosquitoes could help control the disease and limit its spread. Although this is currently not easy, especially as climate change causes global temperatures to slowly rise, it is certainly possible. "Medical researchers may develop some drugs with such an effect that once a mosquito bites a malaria-infected patient who takes the drugs, the EIP of parasites in that mosquito will be prolonged," Wang said. The development of these EIP-extending drugs could lead to a decrease in the spread of malaria and its dire effects.

**More information:** Wang, X., & Zhao, X-Q. A Periodic Vector-Bias Malaria Model with Incubation Period. *SIAM Journal on Applied Mathematics*.

## Provided by Society for Industrial and Applied Mathematics

Citation: Mathematical model limits malaria outbreaks (2017, January 24) retrieved 23 June 2024 from <u>https://phys.org/news/2017-01-mathematical-limits-malaria-outbreaks.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.