

Moth gut bacterium defends its host by making antibiotic

January 19 2017



Cotton leafworm, *Spodoptera littoralis*. Credit: S. Thiessen, MPI for Chemical Ecology

Nearly half of all insects are herbivores, but their diets do not consist of only plant material. It is not uncommon for potentially harmful microorganisms to slip in during a feast. In a study published on January 19 in *Cell Chemical Biology*, researchers report that these insects use an ironic strategy to resist microbial infections. A bacterial species commonly found in the gut of the cotton leafworm and other moths secretes a powerful antimicrobial peptide, killing off competitors while defending its host against pathogens.



"It has long been proposed that native gut bacteria are an important component of host defense, but until now, the responsible <u>species</u> and molecular mechanism have not been clearly demonstrated," says first author Yongqi Shao of Zhejiang University. "We show that the evolutionary success of insects is partially based on a symbiotic association with <u>gut microbes</u>, which co-evolved with their hosts over millions of years."

Insects are the largest group of animals on Earth, with over a million known species. As the most significant herbivores in the world, insects ingest microbes commonly found in plants or the surrounding environment, yet they are remarkably resistant to infections. Increasing evidence in both vertebrates and invertebrates suggests that gut bacteria defend hosts against invading microbes. But the species that exert this protective effect have rarely been identified, leaving the molecular mechanism of action unclear.

A clue to this question came while Shao and senior study author Wilhelm Boland of the Max Planck Institute for Chemical Ecology were studying the cotton leafworm, *Spodoptera littoralis*, which is one of the most widespread insect herbivores in the temperate regions and causes substantial economic losses in crop production. They noticed that the composition of gut microbes colonizing this pest changes dramatically during larval development. Whereas young larvae were inhabited by a variety of virulent *Enterococcus species*, older larvae were dominated by *E. mundtii*, which has rarely been documented as a pathogen.





A group of cotton leafworms, Spodoptera littoralis. Credit: Yongqi Shao

Based on these intriguing observations, the researchers set out to identify the factors that influence the composition of gut microbes during insect development. To do so, they isolated different *Enterococcus* species from cotton leafworm larvae and examined how they compete with one another. They discovered that E. mundtii inhibited the growth of other related bacterial species by secreting an <u>antimicrobial peptide</u> called mundticin KS. While this toxin was undetectable in young larvae, it dramatically increased in abundance in older larvae due to expansion of the *E. mundtii* population, shaping the microbiome with surprising efficiency.



"The antimicrobial promotes symbiosis by providing a competitive advantage for *E. mundtii*, contributing to its dominance in the gut microbiome, while protecting the cotton leafworm against pathogens," Boland says. "We expect that protective associations with antibioticproducing bacteria is a common strategy of insects against microbial invaders."

In future studies, the researchers will examine whether similar mechanisms exist in other insect species and look for additional toxic compounds that shape the microbiome during host development. In the end, the findings could have widespread implications for agriculture and health. For example, antimicrobial peptides could be used as food preservatives, and understanding the role of indigenous gut residents could contribute to the development of novel biocontrol strategies against herbivorous insect pests.

"Our study also provides interesting approaches for medical research," Shao says. "Many conventional antibiotics are facing increasing problems of resistance. As evolutionarily conserved weapons, bacteriocins have great potential as alternatives to conventional antibiotics."

More information: *Cell Chemical Biology*, "Symbiont-derived antimicrobials contribute to the control of the lepidopteran gut microbiota" <u>www.cell.com/cell-chemical-bio ... 2451-9456(16)30438-X</u>, <u>DOI: 10.1016/j.chembiol.2016.11.015</u>

Provided by Cell Press

Citation: Moth gut bacterium defends its host by making antibiotic (2017, January 19) retrieved 2 May 2024 from <u>https://phys.org/news/2017-01-moth-gut-bacterium-defends-host.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.