


Oceanographic analysis offers potential crash site of MH370

January 23 2017



Trajectories of undrogued surface drifters from the historical dataset. The blue trajectories correspond to drifters washing ashore (otherwise the grey colour is used). The small red dots represent the location of their latest transmission. The larger circles refer to drifters reaching land: in red, for the drifters that at some time travelled through the search area; in blue, the same but for the indirect trajectories. The thicker trajectories represent those drifters arriving at Reunion Island. The one in green represents a trajectory that at some point passed through the search area. Credit: J. A. Trinanes et. al.

A group of oceanographers offers a new analysis of the potential crash site of flight Malaysian Airlines flight 370 in the southern Indian Ocean. The researchers, which included scientists from the University of Miami (UM) Rosenstiel School of Marine and Atmospheric Science, used data from buoys that monitor ocean conditions.

In their analysis the team considered the trajectories of drifting buoys, called drifters, from NOAA's Global Drifter Database and of an ocean numerical model. The researchers included only those data from drifters that were unanchored, or undrogued, to better simulate the buoyancy conditions of airplane debris. The team then produced a simulation model of drifter motion using known oceanographic conditions near the potential crash site.

The analysis showed that it would take six months to one year for the drifters to reach western Australia and one-and-a half to two years to reach eastern Africa. Interestingly, two drifters traveled from the search region to the area of Reunion Island during the period between the crash of flight MH370 and when the airplane flaperon was found.

These results are consistent with the time and location of the aircraft debris that was found off Reunion Island, almost 17 months after the plane disappeared, and with the recently confirmed finding in Mozambique almost two years later.

The trajectories of the undrogued drifters and synthetic drifters revealed several areas of high probability in the southern Indian Ocean where debris from the missing flight could have passed, including vast areas of the South Indian Ocean, some of them in the relative neighborhood of the search area.

This study "highlights the importance of sustained observations to monitor [ocean conditions](#) that may serve a suite of applications and

studies," said the authors.

The methods developed by the researchers for use in the study could also help scientists track oil spills, and other types of [marine debris](#) and pollutants in the [ocean](#).

More information: Joaquin A. Trinanes et al, Analysis of flight MH370 potential debris trajectories using ocean observations and numerical model results, *Journal of Operational Oceanography* (2016). [DOI: 10.1080/1755876X.2016.1248149](https://doi.org/10.1080/1755876X.2016.1248149)

Provided by University of Miami

Citation: Oceanographic analysis offers potential crash site of MH370 (2017, January 23) retrieved 9 May 2024 from <https://phys.org/news/2017-01-oceanographic-analysis-potential-site-mh370.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.