

Scientists discover bees prefer warm violets in cool forests

January 6 2017



Bees, especially fuzzy females of Carlin's bee (*Andrena carlinii*), prefer to forage upside down on these flowers so their hind legs and bee butts are warmed by the dark petals as they drink nectar and collect pollen. Credit: Dr. Zong-Xin Ren

Research by scientists at Saint Louis University's Bernhardt/Meier

Laboratory engaged in a study of Missouri bees and wildflowers has been published in the online *Journal of Pollination Ecology*.

Peter Bernhardt, Ph.D., a professor of biology at SLU and research associate at the Missouri Botanical Garden and the Royal Botanic Gardens and Domain Trust in Sydney, New South Wales, has been studying reproductive patterns in wildflowers in six countries for over 40 years and, like most dedicated scientists, thrives on new discoveries such as how bees respond to the color of the flowers they pollinate.

"Remember how you were told that a dark coat keeps you a little warmer on a cold but sunny day," Bernhardt said. "Some plants blooming in chilly environments have dark purple or almost black patches on their flowers to keep cold-blooded insects toasty warm as they pollinate.

Bernhardt said three years of research at their lab, with field work at Missouri's Cuivre River State Park and the Shaw Nature Reserve (owned by the Missouri Botanical Garden) illustrate a new side to this colorful tale in the [online journal](#).

The birds foot violet (*Viola pedata*) has two, common, color forms when it blooms during the cool, Missouri, April. The concolor form makes flowers with five light violet-mauve petals. The flower of a bicolor plant has three mauve petals plus two top petals that are a deep, dark, funereal purple.

Co-researcher Retha Edens-Meier, Ph.D., a professor and research scientist in SLU's School of Education, using thermocouples, and a hypodermic tissue probe, learned that these dark petals are up to 3 degrees Celsius (5.4 degrees Fahrenheit) warmer than the surrounding atmosphere when they stand in a pool of Spring sunlight. Bees, especially fuzzy females of Carlin's bee (*Andrena carlinii*), prefer to forage upside down on these flowers so their hind legs and bee butts are

warmed by the dark petals as they drink nectar and collect pollen.

"What is so unusual about these findings is that, when given a choice over two years, native bees preferred to forage on the concolor form so concolor flowers made more seeds compared to bicolors," said Bernhardt."

Comparing violet populations, at two isolated sites, the research team noted that when the plants grew in a sunny, open, limestone glade (Shaw Nature Reserve) the concolors outnumbered the bicolors by 40 to one. It was very different on the shady forested slope at Cuivre River State Park where bicolors and concolors occurred in almost equal numbers or bicolors outnumbered concolors by almost two-to-one in one season. As pools of light shifted over the course of the day under the trees in the cooler forest, cold-blooded bees visited the bicolors more frequently, possibly because of the warmth provided by the darker petals.

Edens-Meier also says that these studies provide us with a fresh insight into how such tiny wild flowers continue to thrive and reproduce.

"Bird's-foot violets have an unusually broad distribution from eastern Canada through at least 30 American states. Research on these botanical beauties reveals interesting information valuable to ecologists, conservationists, and the general public," Edens-Meier said. "Although bird's-foot violets are not endangered within the United States plenty of other plant species are in significant decline."

"Our goals are to investigate plant breeding systems and pollination ecology, especially in rare and threatened plants. As climate change continues to be responsible for out-of-sync bee emergence with flowering periods and bee-specific pollination events, pollination ecology has become an ever-increasing valuable field of study that helps us understand the ecological impact of these environmental changes.

Through research and education, we can attempt to save our threatened organisms, one species at a time," Edens-Meier added.

More information: BERNHARDT, Peter et al. Comparative floral ecology of bicolor and concolor morphs of *Viola pedata* (Violaceae) following controlled burns. *Journal of Pollination Ecology*, [S.l.], v. 19, nov. 2016. ISSN 1920-7603. www.pollinationecology.org/index.php/article&op=view&path%5B%5D=403&path%5B%5D=136

Provided by Saint Louis University

Citation: Scientists discover bees prefer warm violets in cool forests (2017, January 6) retrieved 1 May 2024 from <https://phys.org/news/2017-01-scientists-bees-violets-cool-forests.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--