

Stalagmites in dry corridor suggest Amazonia maintained forests during the last ice age

January 12 2017, by Bob Yirka



Uncontacted indigenous tribe in the brazilian state of Acre. Credit: Gleilson Miranda / Governo do Acre / Wikipedia

(Phys.org)—An international team of researchers has found evidence that suggests that Amazonia did not revert to savannah during the last ice age and instead remained forested. In their paper published in the journal *Nature*, the researchers describe how they retrieved stalagmite samples from a cave in one of the drier parts of Amazonia and used them to calculate not only how dry the area became during the last ice age, but what sorts of vegetation were growing. Mark Bush with the Florida Institute of Technology offers a News & Views piece in the same journal issue on the work done by the team and further explains how measuring the abundance of oxygen isotopes in stalagmites can reveal so much geological information.

Earth scientists have debated for many years how Amazonia fared during the Last Glacial Maximum (LGM)—some have speculated that it was likely the rainforests disappeared altogether due to less [rainfall](#) and was replaced by savannah. Others have countered that it was possible the forests remained throughout the LGM due to other difficult to discern factors. In this new effort, the researchers claim to have found evidence that supports the latter view.

To learn more about the conditions present during the LGM, the researchers ventured to Paraíso Cave, located in what has been described as the dry corridor in the Amazonian lowlands, due to the area receiving less rainfall than other parts of the rain forest. In the cave, they located and removed seven stalagmites which they brought back to their lab for study. By studying the [oxygen isotopes](#) in them, the researchers were able to see that rainfall in the area (which percolated through the ground into the cave) was approximately half (42 percent) that seen today and that the local landscape remained heavily forested.

The researchers suggest that the reason the forests remained was lower temperatures in the area, which reduced evaporation, offsetting the loss of rainfall. Bush notes that data presented by the researchers studying the

stalagmites is consistent with data from a prior study that examined lake sediments in the area. He also notes that it appears likely that fire in the area during the LGM was rare because it has been known to transform wet forests into savannah.

More information: Xianfeng Wang et al. Hydroclimate changes across the Amazon lowlands over the past 45,000 years, *Nature* (2017). DOI: [10.1038/nature20787](https://doi.org/10.1038/nature20787)

Abstract

Reconstructing the history of tropical hydroclimates has been difficult, particularly for the Amazon basin—one of Earth's major centres of deep atmospheric convection^{1, 2}. For example, whether the Amazon basin was substantially drier^{3, 4} or remained wet^{1, 5} during glacial times has been controversial, largely because most study sites have been located on the periphery of the basin, and because interpretations can be complicated by sediment preservation, uncertainties in chronology, and topographical setting⁶. Here we show that rainfall in the basin responds closely to changes in glacial boundary conditions in terms of temperature and atmospheric concentrations of carbon dioxide⁷. Our results are based on a decadal resolved, uranium/thorium-dated, oxygen isotopic record for much of the past 45,000 years, obtained using speleothems from Paraíso Cave in eastern Amazonia; we interpret the record as being broadly related to precipitation. Relative to modern levels, precipitation in the region was about 58% during the Last Glacial Maximum (around 21,000 years ago) and 142% during the mid-Holocene epoch (about 6,000 years ago). We find that, as compared with cave records from the western edge of the lowlands, the Amazon was widely drier during the last glacial period, with much less recycling of water and probably reduced plant transpiration, although the rainforest persisted throughout this time.

© 2017 Phys.org

Citation: Stalagmites in dry corridor suggest Amazonia maintained forests during the last ice age (2017, January 12) retrieved 2 May 2024 from <https://phys.org/news/2017-01-stalagmites-corridor-amazonia-forests-ice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.