


# Topological photonic crystal made of silicon

January 19 2017



(Top) Schematic of photonic crystals consisting of nanorods derived from the honeycomb lattice viewed from above, and (bottom) the corresponding photonic bands. Photonic crystals are obtained by dividing the nearest neighboring nanorods into hexagonal clusters, and widening (a) or narrowing (c) the separation between the hexagonal clusters from the original honeycomb lattice (b), while maintaining the same shape and size of the hexagons. In (c), a band inversion occurs between the photonic  $p$ - and  $d$ -bands, which generates topological features in the system. Credit: National Institute for Materials Science

WPI-MANA researchers derive topological photonic states purely based

on silicon, which can lead to the development of new functions and devices through integration with semiconductor electronics

Topology is a mathematical concept which describes the way of connection of an object invariant under continuous deformation. Recently, it has been pointed out that topology can also be defined in the electronic states of [materials](#), and this provides a unified and useful picture for physicists describing the unique properties of materials.

Based on a new approach which they have dubbed "topological nanoarchitectonics," Xiao Hu and Long-Hua Wu, who are theoreticians at the International Center for Materials Nanoarchitectonics (MANA), National Institute for Materials Science (NIMS), elucidated a new principle which makes [electromagnetic waves](#) including light propagate on the edge in a two-dimensional photonic crystal without being scattered.

It was known that scattering of light by defects in conventional photonic crystals can be suppressed in topological photonic states, but so far special materials were required in order to create topological photonic crystals. However, the MANA researchers discovered a new principle which makes it possible to realize a topological photonic crystal by merely adjusting the positions of insulator or semiconductor nanorods in a [honeycomb lattice](#), without using any special material or complicated structure. When hexagonal clusters are formed by adjusting the positions of nanorods, electromagnetic modes carrying spin, which is conventionally specific to electrons, appear. As a result, the MANA researchers theoretically clarified that a photonic crystal exhibits topological properties when the separation between hexagonal clusters is narrowed from that of the honeycomb lattice.

Since this new property of a [photonic crystal](#) can be obtained even by semiconductor such as silicone and/or GaN alone, various new functions

are expected by integration of information processing functions achieved by the well-established semiconductor electronics and the excellent topological property of electromagnetic waves.

**More information:** Long-Hua Wu et al. Scheme for Achieving a Topological Photonic Crystal by Using Dielectric Material, *Physical Review Letters* (2015). [DOI: 10.1103/PhysRevLett.114.223901](https://doi.org/10.1103/PhysRevLett.114.223901)

Provided by National Institute for Materials Science

Citation: Topological photonic crystal made of silicon (2017, January 19) retrieved 18 April 2024 from <https://phys.org/news/2017-01-topological-photonic-crystal-silicon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.