


# Biologists report findings on ants' internal navigation systems

February 16 2017

---



Desert ant in the spherical treadmill. Credit: Matthias Wittlinger

Biologists of the University of Freiburg have used a spherical treadmill to investigate how desert ants navigate in a featureless environment. Cataglyphis desert ants live in salt pans and are ideal models for such

navigation research. When they set out in search of food in their flat, bare, hostile environment, they are always able to find their way back to their nest via the shortest route possible using an internal navigation system. The biologists published their results in the *Journal of Experimental Biology*.

The ants measure the distance they have traveled by recording how many steps they have taken—and they use the sun for directional orientation, taking into account its movement over time via their own internal clock. A team of researchers led by Dr. Matthias Wittlinger of the University of Freiburg developed a tiny treadmill on which the ants behave just as they do in the wild. "This gives us almost unlimited possibilities to test the mechanisms and neural basis of our model animal's spatial orientation and navigation in the laboratory," says Wittlinger. "We can place the ants in a virtual world and incorporate certain changes to see how they react." The experiments are expected to yield information useful in the development of autonomous robots, as well as in other areas.

The ant treadmill is like a ball on top of which the insect can walk, a bit like a hamster in a wheel. To the ant, it is like walking in its normal environment—although it doesn't really go anywhere. The team developed the spherical treadmill in such a way that it enables the ant to walk using a natural gait, even when moving and changing direction quickly. The spherical treadmill contains optical sensors like those in a computer mouse to record precisely the ant's direction and speed.

In the experiments, the ant initially travels some ten meters from its nest—whether in the field or in the laboratory—storing information about the path it has taken. Then the researchers place it into the treadmill. The ant then moves exactly as it would if it were returning to its nest. It first heads back to the nest as directly as possible. Once it has arrived in the general area of the nest, the ant switches to search mode,

taking a meandering path so as to find the exact location. The biologists have observed that the [ants](#) adapt their speed to the relevant phase of their journey—fast on the initial homeward trip, then slow in the searching phase. Comparisons between the virtual journey inside the treadmill and the necessary trip on the ground show that the insects have a high-precision navigation system.

**More information:** Hansjuergen Dahmen, Verena Luisa Wahl, Sarah Elisabeth Pfeffer, Hanspeter Mallot, and Matthias Wittlinger (2017): Naturalistic path integration of *Cataglyphis* desert ants on an air cushioned light-weight spherical treadmill. In: *Journal of Experimental Biology* 220/4.

Provided by University of Freiburg

Citation: Biologists report findings on ants' internal navigation systems (2017, February 16) retrieved 10 April 2024 from <https://phys.org/news/2017-02-biologists-ants-internal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.