

Researchers discover a black hole feeding frenzy that breaks records

February 6 2017



Artist illustration depicting the record breaking "tidal disruption event" (TDE). The red shows hotter material that falls toward the black hole and generates a distinct X-ray flare. The blue shows a wind blowing from the infalling material. Credit: CXC/M. Weiss; X-ray: NASA/CXC/UNH/D. Lin et al, Optical: CFHT



A giant black hole ripped apart a nearby star and then continued to feed off its remains for close to a decade, according to research led by the University of New Hampshire. This black hole meal is more than 10 times longer than any other previous episode of a star's death.

"We have witnessed a star's spectacular and prolonged demise," said Dacheng Lin, a research scientist at UNH's Space Science Center and the study's lead author. "Dozens of these so-called <u>tidal disruption</u> events have been detected since the 1990s, but none that remained bright for nearly as long as this one."

Using data from a trio of orbiting X-ray telescopes, NASA's Chandra Xray Observatory and Swift Satellite as well as ESA's XMM-Newton, researchers found evidence of a massive "tidal disruption event" (TDE). Tidal forces, due to the intense gravity gradient of the black hole, can destroy an object – such as a star – that wanders too close. During a TDE, some of the stellar debris is flung outward at high speeds, while the rest falls toward the black hole. As it travels inward, and is ingested by the black hole, the material heats up to millions of degrees and generates a distinct X-ray flare.

These multiwavelength flares, which can be viewed by the satellites, help to study otherwise dormant massive back holes. Previous flares were short-lived, typically becoming very faint in a year, but this super-long Xray flare has been persistently bright for close to a decade. The extraordinary long bright phase of this TDE means that either this was the most massive star ever to be torn apart during one of these events, or the first where a smaller star was completely torn apart.

The X-ray source containing this force-fed black hole, known by its abbreviated name of XJ1500+0154, is located in a small galaxy about 1.8 billion light years from Earth.



The X-ray data also indicates that radiation from material surrounding this black hole has consistently surpassed the so-called Eddington limit, defined by a balance between the outward pressure of radiation from the hot gas and the inward pull of the gravity of the black hole.

The conclusion that <u>supermassive black holes</u> can grow, from TDEs and perhaps other means, at rates above those corresponding to the Eddington limit has important implications. Such rapid growth may help explain how supermassive <u>black holes</u> were able to reach masses about a billion times higher than the sun when the universe was only about a billion years old.

Based on the modeling by the researchers the black hole's feeding supply should be significantly reduced in the next decade and begin to fade in the next several years.

A paper describing these results appears in the February 6th issue of the journal *Nature Astronomy*.

More information: A likely decade-long sustained tidal disruption event, *Nature Astronomy*, <u>www.nature.com/articles/s41550-016-0033</u>, <u>arxiv.org/abs/1702.00792</u>

Provided by University of New Hampshire

Citation: Researchers discover a black hole feeding frenzy that breaks records (2017, February 6) retrieved 2 May 2024 from <u>https://phys.org/news/2017-02-black-hole-frenzy.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.