

# Great Barrier Reef building coral under threat from poisonous seaweed

February 3 2017

---



Credit: Griffith University

The Griffith University study, conducted in collaboration with national

and international experts in reef and chemical ecology, showed that if the world continues with 'business as usual' CO<sub>2</sub> emissions important reef building corals will suffer significantly by 2050 and die off by 2100.

Associate Professor Guillermo Diaz-Pulido, of Griffith's School of Environment, says that is because algae will compete for space with corals in the reef, much like a weed, and eventually take over.

Researchers knew increased CO<sub>2</sub> had an effect on seaweed behaviour but have now been able to demonstrate just how this happens. They discovered this is due to an increase in the potency of chemical compounds that poison corals.

"This is a major step forward in understanding how seaweeds can harm corals and has important implications for comprehending the consequences of increased carbon dioxide emissions on the health of the Great Barrier Reef," says Associate Professor Diaz-Pulido.

"For the algae to grow they need light and CO<sub>2</sub>, just like any other plant, and because algae in the future would be exposed to much more CO<sub>2</sub> in seawater we wanted to know to what extent the CO<sub>2</sub> would affect some of the things algae do, the physiology and the interaction with animals."

Professor Mark Hay, from the Georgia Institute of Technology and co-author of the study, adds: "What we've discovered is that some algae produce more potent chemicals that suppress or kill corals more rapidly. This can occur rapidly, in a matter of only weeks.

"If the algae overtake the coral we have a problem which contributes to reef degradation, on top of what we already know with coral bleaching, crown of thorn starfish outbreaks, cyclones or any other disturbance."

The research was undertaken at Heron Island, a [coral](#) cay on the southern end of the reef using underwater reef experiments and outdoor lab studies.

Associate Professor Diaz-Pulido says the study has global impacts because one of the seaweeds studied that causes the most damage is a common brown alga species found in reefs worldwide.

"That's a problem because if these [algae](#) take advantage of elevated CO<sub>2</sub> in seawater that's even more a matter of concern," he says.

"The scale of the problem is so big removing a bunch of seaweed from the [reef](#) isn't going to do much because it just regrows and regenerates, so I think the way to address this really is to reduce the levels of CO<sub>2</sub> in the atmosphere."

**More information:** Carlos Del Monaco et al, Effects of ocean acidification on the potency of macroalgal allelopathy to a common coral, *Scientific Reports* (2017). [DOI: 10.1038/srep41053](https://doi.org/10.1038/srep41053)

Provided by Griffith University

Citation: Great Barrier Reef building coral under threat from poisonous seaweed (2017, February 3) retrieved 28 April 2024 from <https://phys.org/news/2017-02-great-barrier-reef-coral-threat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---