

Radar study targeting wind events at NASA, Air Force launch facilities

February 10 2017, by Phillip Gentry

Rockets rolling to their launch towers at both the Cape Canaveral Air Force Station and NASA's Kennedy Space Center in Florida (and the people who work on them) may soon be a little bit safer, thanks to research in the Atmospheric Science Department at The University of Alabama in Huntsville (UAH).

Funded by NASA's Marshall Space Flight Center and the U.S. Air Force's 45th Weather Squadron (45WS), research by UAH graduate student Corey Amiot has focused on improving lead times for warnings of high [wind](#) events that threaten the Florida launch complexes.

Now, Amiot's task is to verify the extended warning times (up to 40 minutes in advance of a high speed wind event) and then to figure out how to reduce the number of false alarms.

"I'm looking at thunderstorm downbursts, when a convective storm begins to collapse and spread out at the surface," Amiot said.

His primary tool is storm data collected by the dual-polarization C-band weather radar operated by the 45WS from its station 26.5 miles southwest of the launch facilities. He also uses data from 29 weather stations scattered across the NASA and Air Force facilities.

"I have those towers as ground truth," Amiot said, "and a way to identify which storms produced wind gusts on the ground."

Early results from this research were presented recently at an American Meteorological Society meeting in Seattle.

NASA and the Air Force want Amiot to improve predictions of ground wind in excess of 35 knots (just over 40 mph), the point at which strong wind warnings are issued and personnel are required to take safety measures. Wind that speed or faster could be hazardous to people working outdoors and might put some hardware at risk.

Amiot used the dual-polarization radar data from 14 high-wind producing warm-season thunderstorms to look for patterns and possible signatures of upcoming changes in each storm. Dual-polarization radar sends out both vertical and horizontal radar waves. By comparing the signals returned by both, researchers can look inside a storm and see water droplets, small ice particles and larger ice that might indicate the presence of graupel or hail.

"A storm updraft lifts liquid water above the freezing level, where it forms ice," Amiot said. "This is very important to downburst formation. When the ice melts it cools the surrounding air, which will accelerate the downburst."

Focusing on multi-cell storms, Amiot identified four distinct radar signatures found in 85 to 92 percent of the 14 storms that generated high wind events.

"This is a good step forward," he said. "Now I need to expand the sample size so I can identify other [radar](#) signatures in addition to the four I've identified so far."

Provided by University of Alabama in Huntsville

Citation: Radar study targeting wind events at NASA, Air Force launch facilities (2017, February 10) retrieved 19 April 2024 from <https://phys.org/news/2017-02-radar-events-nasa-air-facilities.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.