

# Novel analytical techniques to detect solar radiation imprints on meteoroids

March 22 2017, by Barbara Vonarburg

---




The meteorites being studied were collected in Oman by a team coordinated by Beda Hofmann from the Natural History Museum Bern. Credit: Hofmann / Natural History Museum Bern

When a meteoroid travels in space, solar radiation leaves distinctive imprints on its outer layer. Together with colleagues, ETH researcher Antoine Roth has developed novel analytical techniques to detect these imprints, allowing the team to reconstruct meteorites' space journeys.

The inconspicuous, small stone that was analysed with high-tech equipment is named Jiddat al Harasis 466. It travelled a long way before it entered the Earth's atmosphere and landed in the desert of Oman. "We think that Jiddat al Harasis 466 was formed 4 million years ago as a

remnant of a crash of bigger blocks in the [asteroid belt](#) between Mars and Jupiter," explains Antoine Roth from the Institute of Geochemistry and Petrology at ETH Zurich. Then, after a rapid transfer from the asteroid belt to Earth, it heated up intensely during atmospheric entry and lost a lot of material. What was originally a stone with a radius of two centimetres ended up as a one centimetre [meteorite](#).

Jiddat al Harasis 466 is one of 25 small meteorites that Roth chose for his study, which is now to be published by the journal *Meteoritics & Planetary Science*. In order to find out more about the history of the samples, he was looking for [neon](#): a [noble gas](#) that can be produced by energetic solar radiation, for instance when it splits the magnesium atoms that are part of the rocky material. Knowing the amount of neon produced by solar cosmic ray allows scientists to figure out how far from the sun and how long a meteorite travelled in space. "It's like being able to tell whether your friends spent their holiday at a sunny beach or in a cold place because of their tan," says Roth, who is also a member of the Swiss National Centre of Competence in Research PlanetS.



Photomicrograph of Jiddat al Harasis 466 (thin-section). Credit: A.Roth / ETH Zurich

Previously, neon produced by solar cosmic rays had been found in Martian meteorites, but not in ordinary "chondrites" originating from the asteroid belt. "This may be the result of a sampling bias," Roth says, "because neon produced by solar radiation is better preserved in meteorites with small pre-atmospheric radii – and these specimens are often studied only if they belong to unusual or rare classes." Since the solar cosmic rays penetrate only a few centimetres into the rocky material, with bigger samples the neon gets lost when the meteorite is ablated during its entry into Earth's atmosphere. But in small samples,

the noble gas can be preserved in the centre.

## Samples from Omani–Swiss project

Looking for small meteorites, the scientist found a rich collection at the Natural History Museum in Bern, which coordinated several meteorite search campaigns in Oman. Using an infrared laser and a mass spectrometer at the University of Bern, the researchers were able to extract the neon from the samples and measure its isotopic concentrations: this allows them to determine the proportion of the noble gas that originated from solar cosmic rays rather than [galactic cosmic rays](#). To analyse the measured data and calculate the actual production rate, Roth and his colleagues developed a new physical model that also predicts the average distance to the sun at which the meteoroid has been irradiated.

As a result, Roth found neon produced by [solar radiation](#) in 4 of the 25 chondrites that were studied. Some of the samples that didn't show the sought-after noble gas were probably part of a bigger stone that first fell apart during entry into Earth's atmosphere. "Our data indicate that neon produced by solar cosmic rays is by no means limited to Martian meteorites," summarises the PlanetS member. In an upcoming study he will analyse meteorites collected by NASA in Antarctica. Since he needs small samples that weigh less than 10 grams each, it was not too complicated to get appropriate material – although the chondrites are destroyed during the analysis.

**More information:** Neon produced by solar cosmic rays in ordinary chondrites. *Meteoritics & Planetary Science*. DOI: 10.1111/maps.12868

Provided by ETH Zurich

Citation: Novel analytical techniques to detect solar radiation imprints on meteoroids (2017, March 22) retrieved 1 May 2024 from <https://phys.org/news/2017-03-analytical-techniques-solar-imprints-meteoroids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.