

## Computer model shows breakup of iceberg logjams

March 2 2017, by Robert Perkins



Credit: B. Minchew/Caltech

Projections of how much the melting of ice sheets will contribute to sealevel rise can vary by several meters based on the rate of iceberg calving at the edges of those ice sheets. To provide climate scientists with models that make more accurate forecasts, a postdoctoral researcher at Caltech has created a computer simulation of one of the key processes controlling glacial calving.

Glaciers are moving slabs of ice that slowly grind downhill. Where they



end in the sea, chunks break off, forming icebergs in a process known as calving. When temperatures plummet in the winter, those icebergs can freeze together and create a traffic jam that prevents further icebergs from breaking off from the glacier.

During the winter, the glacier loses much less ice to the sea. The eventual spring breakup of what is known as the mélange—that frozen iceberg logjam—occurs suddenly, and is the focus of research by Caltech's Alexander Robel.

"I developed a computer model that simulates how the first iceberg calving of the warm season creates a shock wave that travels through the jammed mélange, breaking it up," says Robel, a National Oceanic and Atmospheric Administration (NOAA) Postdoctoral Scholar and a Stanback Postdoctoral Scholar at Caltech. His new model was featured in *Nature Communications* on February 28.

The mélange is a frozen granular material, so Robel adapted an opensource computer simulation called the Discrete-Element Bonded-Particle Sea Ice Model to show how icebergs freeze together in the winter and then transmit the shock of the first iceberg calving in the summer.





This animation shows how a single calving event sends out a shockwave that triggers a breakup of the frozen mélange. Credit: Alexander Robel

That first calving is made possible by the thinning of sea ice in warmer water, which reduces the ability of the mélange to act a bulwark against the glacier.

Robel tailored his modeled glaciers to resemble fjords in Greenland. Those fjords are narrow channels of water that are prone to trapping mélange. Robel was able show that the threshold at which spring sea-ice breakup is likely to occur is based in part on the thickness of sea ice within the mélange, but also on the shape of the channel within which the mélange is trapped.

Robel, who is a researcher in Caltech's Division of Geological and Planetary Sciences, home to the Seismological Laboratory, says his work was inspired in part by seismological studies of the way fractures propagate through elastic materials—drawing a connection between earthquakes and <u>iceberg calving</u>.





This animation shows how mélange responds to pressure applied by the movement of a glacier. Credit: Alexander Robel

Robel's paper is titled "Thinning Sea Ice Weakens Buttressing Force of Iceberg Mélange and Promotes Calving."

**More information:** Alexander A. Robel. Thinning sea ice weakens buttressing force of iceberg mélange and promotes calving, *Nature Communications* (2017). DOI: 10.1038/ncomms14596

Provided by California Institute of Technology

Citation: Computer model shows breakup of iceberg logjams (2017, March 2) retrieved 23 June 2024 from <u>https://phys.org/news/2017-03-breakup-iceberg-logjams.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.