

The energetic cost of swimming at high speed when startled may be a factor in strandings of dolphins and whales

March 16 2017, by Kathryn Knight





Dolphins and other cetaceans have to balance their energy demands during dives with a limited oxygen reserve. Credit: T. M. Williams

Dolphins swimming at top speed use more than twice the amount of energy per fin beat than dolphins swimming at a more relaxed pace, according to a study by scientists at UC Santa Cruz. The researchers also found that startled beaked whales fleeing human noises use 30.5 percent more energy during the flight, suggesting that the high cost of escape could contribute to recent dolphin and whale strandings.

Diving marine mammals must balance speed and the duration of breathhold with the need to conserve limited oxygen reserves, leaving them vulnerable when they try to escape from perceived threats. "Amazingly, there has been only a handful of studies that have actually measured the energetic cost of a dive for <u>dolphins</u> or whales," said Terrie Williams, professor of ecology and evolutionary biology at UC Santa Cruz.

Williams is fascinated by how marine mammals balance their energy demands with their finite oxygen supply. Explaining that fleeing dolphins beat their fins continually when <u>swimming</u> full-out, while adopting a more leisurely burst-and-glide style during a routine dive, Williams said she wondered just how much energy each swimming style uses and how much energy a startled animal might use when evading peril.

In a paper published March 15 in the *Journal of Experimental Biology*, Williams and her colleagues reported the energetic costs of different swimming strokes for trained dolphins and other cetaceans. Using that data and previously reported observations of wild Cuvier's <u>beaked</u> <u>whales</u> fleeing from low-frequency sonar, they were able to calculate



how much more energy the beaked whales expended in their efforts to escape the noise.



Williams and her colleagues trained bottlenose dolphins to participate in tests that would allow the scientists to measure the metabolic costs of different swimming styles. Credit: University of California - Santa Cruz

Working with a team of expert trainers, Williams and her colleagues spent over 6 months training six <u>bottlenose dolphins</u> that had previously worked with the U.S. Navy to participate in swimming tests that would allow the scientists to measure the metabolic costs of the different swimming styles. In the first test, the dolphins learned to swim at their most comfortable speed while pushing against a force plate in the wall of the pool as the researchers filmed the number of fin beats. The second test required the animals to dive down 10 meters wearing a fin-beat tracker and swim through a series of hoops before returning to the surface.



Williams was able to take advantage of the animals' marine lifestyle to directly measure the metabolic cost of each dive by training the animals to surface in an air dome where she could record how much oxygen the animals inhaled as they recharged the oxygen stores that they had consumed while swimming. When Williams included killer whales in the metabolic measurements, she had to build an outsized 1.7-square-meter respiration dome to accommodate the larger animals.

After months of patience, the team was eventually able to calculate that bottlenose dolphins consume 3.3 Joules per kilogram per stroke during routine swimming, but the energy consumption almost doubles to 6.4 J/(kg stroke) when swimming their hardest. When the team added the killer whales' fin-beat cost to a plot including the swimming costs of bottlenose dolphins, harbor porpoises and beluga whales, they finally had a tool that they could use to estimate the diving costs of any cetacean.

But what are the conservation implications of the increased cost of each fin beat when whales and dolphins need to avoid danger? Loud manmade noise pollution is thought to be responsible for some mass strandings, so Williams contacted Brandon Southall, a research associate at UCSC's Institute of Marine Sciences, who had recorded how a Cuvier's beaked whale reacted to 20 minutes of loud sonar. With the recording showing that the whale's fin-beat pattern increased significantly, from about 13.6 to 16.9 strokes per minute, she calculated that the startled animals would use 30.5 percent more energy as their metabolic rate rocketed to power the fleeing animals' fin beats. And the whale did not recover swiftly, continuing to use the most costly fin beats for almost two hours after the noise stopped.

"Not all strokes are the same in terms of energy expenditure for swimming dolphins, and this has enormous implications for the cost of flight from aversive stimuli by wild cetaceans," Williams said. "In view of the number of cetacean mass strandings across the globe and the



increase in human presence in the oceans, such data are critical. The animals in our care provided that opportunity."

More information: Kathryn Knight. High-speed dolphins burn double calories, *The Journal of Experimental Biology* (2017). DOI: 10.1242/jeb.158683

Provided by University of California - Santa Cruz

Citation: The energetic cost of swimming at high speed when startled may be a factor in strandings of dolphins and whales (2017, March 16) retrieved 3 May 2024 from <u>https://phys.org/news/2017-03-energetic-high-startled-factor-strandings.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.