


# How a mineral found in Martian meteorites may provide clues to ancient abundance of water

March 6 2017, by Glenn Roberts Jr.

---



An impact crater on Mars, named Melas Dorsa, and its surroundings show a rich geologic history. The image was created by the European Space Agency's Mars Express. Studies of the transformation of a synthetic version of a mineral known as whitlockite suggest that Mars had a more water-rich past than previously thought. Credit: G. Neukum/ESA,DLR, FU Berlin

Mars may have been a wetter place than previously thought, according to

research on simulated Martian meteorites conducted, in part, at the Department of Energy's Lawrence Berkeley National Laboratory (Berkeley Lab).

In a study published today in the journal *Nature Communications*, researchers found evidence that a mineral found in Martian meteorites—which had been considered as proof of an ancient dry environment on Mars—may have originally been a hydrogen-containing mineral that could indicate a more water-rich history for the Red Planet.

Scientists at the University of Nevada, Las Vegas (UNLV), who led an international research team in the study, created a synthetic version of a hydrogen-containing mineral known as whitlockite.

## **After shock-compression experiments on whitlockite samples that simulated**

the conditions of ejecting meteorites from Mars, the researchers studied their microscopic makeup with X-ray experiments at Berkeley Lab's Advanced Light Source (ALS) and at Argonne National Laboratory's Advanced Photon Source (APS).

The X-ray experiments showed that whitlockite can become dehydrated from such shocks, forming merrillite, a mineral that is commonly found in Martian meteorites but does not occur naturally on Earth.





Naturally formed crystals of the mineral whitlockite, which is rare on Earth, are visible in this sample on display at Canada's Royal Ontario Museum. Credit: Wikimedia Commons

"This is important for deducing how much water could have been on Mars, and whether the water was from Mars itself rather than comets or meteorites," said Martin Kunz, a staff scientist at Berkeley Lab's ALS who participated in X-ray studies of the shocked whitlockite samples.

"If even a part of merrillite had been whitlockite before, it changes the water budget of Mars dramatically," said Oliver Tschauner, a professor of research in the Department of Geoscience at UNLV who co-led the study with Christopher Adcock, an assistant research professor at UNLV.

And because whitlockite can be dissolved in water and contains phosphorous, an essential element for life on Earth—and merrillite appears to be common to many Martian meteorites—the study could also have implications for the possibility of life on Mars.

"The overarching question here is about water on Mars and its early history on Mars: Had there ever been an environment that enabled a generation of life on Mars?" Tschauner said.

The pressures and temperatures generated in the shock experiments, while comparable to those of a meteorite impact, lasted for only about 100 billionths of a second, or about one-tenth to one-hundredth as long as an actual meteorite impact.


The fact that experiments showed even partial conversion to merrillite in these lab-created conditions, a longer duration impact would likely have produced "almost full conversion" to merrillite, Tschauner said.

He added that this latest study appears to be one of the first of its kind to detail the shock effects on synthetic whitlockite, which is rare on Earth.

Researchers blasted the synthetic whitlockite samples with metal plates fired from a gas-pressurized gun at speeds of up to about half a mile per second, or about 1,678 miles per hour, and at pressures of up to about 363,000 times greater than the air pressure in a basketball.

"You need a very severe impact to accelerate material fast enough to escape the gravitational pull of Mars," Tschauner said.

At Berkeley Lab's ALS, researchers used an X-ray beam to study the microscopic structure of shocked whitlockite samples in a technique known as X-ray diffraction. The technique allowed researchers to differentiate between merrillite and whitlockite in the shocked samples.



The Mojave Crater on Mars, which is believed to be the source of some Martian meteorites found on Earth, is pictured here in a rendering produced by the HIRISE camera on NASA's Mars Reconnaissance Orbiter. Credit: NASA JPL-Caltech, University of Arizona

Separate X-ray experiments carried out at Argonne Lab's APS showed that up to 36 percent of whitlockite was transformed to merrillite at the site of the metal plate's impact with the mineral, and that shock-generated heating rather than compression may play the biggest role in whitlockite's transformation into merrillite.


There is also evidence that liquid water flows on Mars today, though there has not yet been scientific proof that life has ever existed on Mars. In 2013, planetary scientists reported that darkish streaks that appear on Martian slopes are likely related to periodic flows of water resulting from changing temperatures. They based their analysis on data from

## NASA's Mars Reconnaissance Orbiter.

And in November 2016, NASA scientists reported that a large underground body of water ice in one region of Mars contains the equivalent of all of the water in Lake Superior, the largest of the Great Lakes. Rover explorations have also found evidence of the former abundance of water based on analysis of surface rocks.

"The only missing link now is to prove that (merrillite) had, in fact, really been Martian whitlockite before," Tschauner said. "We have to go back to the real meteorites and see if there had been traces of water."

Adcock and Tschauner are pursuing another round of studies using infrared light at the ALS to study actual Martian meteorite samples, and are also planning X-ray studies of these actual samples this year.



The dark, narrow, 100-meter-long streaks flowing downhill within the Hale Crater on Mars in this rendering are inferred to have been formed by flowing

water in this false-color image that was draped on a digital terrain model. Credit: NASA JPL, University of Arizona

Many Martian meteorites found on Earth seem to come from a period of about 150 million to 586 million years ago, and most are likely from the same region of Mars. These meteorites are essentially excavated from a depth of about a kilometer below the surface by the initial impact that sent them out into space, so they aren't representative of the more recent geology at the surface of Mars, Tschauer explained.

"Most of them are very similar in the rock composition as well as the minerals that are occurring, and have a similar impact age," he said. Mars is likely to have formed about 4.6 billion years ago, about the same time as Earth and the rest of our solar system.

Even with more detailed studies of Martian meteorites coupled with thermal imaging of Mars taken from orbiters, and rock samples analyzed by rovers traversing the planet's surface, the best evidence of Mars' water history would be an actual Martian rock taken from the planet and transported back to Earth, intact, for detailed studies, researchers noted.

"It's really important to get a rock that hasn't been 'kicked'" like the Martian meteorites have, said Kunz, in order to learn more about the planet's [water](#) history.

Provided by Lawrence Berkeley National Laboratory

Citation: How a mineral found in Martian meteorites may provide clues to ancient abundance of water (2017, March 6) retrieved 18 April 2024 from <https://phys.org/news/2017-03-mineral-martian-meteorites-clues-ancient.html>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.