

NASA spacecraft investigate clues in radiation belts

March 27 2017, by Mara Johnson-Groh



The twin Van Allen Probes orbit one behind the other, investigating clues in a



way a single spacecraft never could. In this model, the trailing spacecraft saw an increase in injected oxygen particles (blue), which was unobserved by the first. The increase in particles was due to a geomagnetic storm front that moved across the path of the orbit after the first spacecraft passed. Credit: NASA's Goddard Space Flight Center/Mike Henderson/Joy Ng, Producer

High above Earth, two giant rings of energetic particles trapped by the planet's magnetic field create a dynamic and harsh environment that holds many mysteries—and can affect spacecraft traveling around Earth. NASA's Van Allen Probes act as space detectives, to help study the complex particle interactions that occur in these rings, known as the Van Allen radiation belts. Recently, the spacecraft were in just the right place, at just the right time, to catch an event caused by the fallout of a geomagnetic storm as it happened. They spotted a sudden rise in particles zooming in from the far side of the planet, improving our understanding of how particles travel in near-Earth space.

The two twin Van Allen Probe spacecraft orbit one behind the other, investigating clues in a way a single spacecraft never could. On one typical day, as the first instrument traveled around Earth, it spotted nothing unusual, but the second, following just an hour later, observed an increase in oxygen particles speeding around Earth's dayside—the side nearest the sun. Where did these particles come from? How had they become so energized?

Scientists scoured the clues to figure out what was happening. With the help of computer models, they deduced that the particles had originated on the night side of Earth before being energized and accelerated through interactions with Earth's <u>magnetic field</u>. As the particles journeyed around Earth, the lighter hydrogen <u>particles</u> were lost in collisions with the atmosphere, leaving an oxygen-rich plasma. The



findings were presented in a recent paper in Geophysical Review Letters.

The unique double observations of the Van Allen Probes help untangle the complex workings of Earth's magnetic environment. Such information has provided the very first view of these harsh belts from the inside—and it helps us better protect satellites and astronauts traveling through the region.

More information: M. H. Denton et al. The complex nature of stormtime ion dynamics: Transport and local acceleration, *Geophysical Research Letters* (2016). DOI: 10.1002/2016GL070878

Provided by NASA's Goddard Space Flight Center

Citation: NASA spacecraft investigate clues in radiation belts (2017, March 27) retrieved 3 May 2024 from <u>https://phys.org/news/2017-03-nasa-spacecraft-clues-belts.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.