

# Nitrogen uptake between fungi and orchids

March 7 2017

---



Researchers investigated *Tulasnella calospora* as both a free-living mycelium and in symbiosis with the photosynthetic orchid long-lipped serapias. Credit: Ziegler175, Wikimedia Commons, CC BY-SA 3.0

Orchids are an example of an experimentally tractable plant that is highly dependent on its relationship with its mycorrhizal fungal partners for nutrient supply. In this recent study, researchers for the first time identified some genetic determinants potentially involved in nitrogen uptake and transfer in orchid mycorrhizas.

This study provides a model system amenable to experimental

manipulation, for plant-fungi nutrient exchanges on a symbiotic level and offers insights into how host plants benefit from the mutualistic relationships formed with [soil fungi](#) which can expand their habitat range. Understanding these vital relationships may shed light on microbial symbioses applicable to growing bioenergy feedstock plants.

Orchids, like the majority of terrestrial plants, form symbiotic relationships between their plant roots and soil fungi, known as mycorrhizal association. However, unlike other terrestrial plants, orchids rely on their mycorrhizal fungal partners for nutrient supply during the feed germination and development stages. Following this stage, most orchid species develop leaves and are capable of self-nourishment whereas some species continue to rely on their fungal partners for an organic carbon supply. In this study, published in the January 2017 issue of *New Phytologist*, a team led by University of Turin researchers investigated the orchid mycorrhizal fungus *Tulasnella calospora* as both a free-living mycelium and in symbiosis with the photosynthetic orchid long-lipped serapias, or *Serapias vomeracea*. For the first time, researchers looked at the fungal genes that may have been involved in both the uptake and transfer of nitrogen to the host plant. RNA sequencing for the project was performed at the U.S. Department of Energy Joint Genome Institute (DOE JGI), a DOE Office of Science User Facility, as part of the 2013 Community Science Program portfolio.

The team also used the DOE JGI fungal genome database MycoCosm to identify [fungal genes](#) coding for proteins that were involved in nitrogen uptake and transfer. They found that the *Tulasnella calospora* genome has two genes coding for ammonium transporters and several genes coding for amino acid transporters, proteins that play roles in the nitrogen nutrient pathway. Overall, the orchid mycorrhizal fungi's use of nitrogen may broaden the habitat ranges of orchids, allowing them to grow in a variety of soil types. Of more general interest to the DOE, this

study provides important insights for this process, and furthers understanding of plant-microbial symbioses that are vital for plant health and may inform understanding of microbial symbioses relevant to bioenergy feedstock plants.

**More information:** Valeria Fochi et al. Fungal and plant gene expression in the-symbiosis provides clues about nitrogen pathways in orchid mycorrhizas, *New Phytologist* (2017). [DOI: 10.1111/nph.14279](https://doi.org/10.1111/nph.14279)

Provided by DOE/Joint Genome Institute

Citation: Nitrogen uptake between fungi and orchids (2017, March 7) retrieved 20 April 2024 from <https://phys.org/news/2017-03-nitrogen-uptake-fungi-orchids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.