




# **Paleolake deposits on Mars might look like sediments in Indonesia**

March 6 2017

---



Lake Towuti, Indonesia. (A) Regional context for Lake Towuti on the island of Sulawesi in Indonesia. Red box indicates approximate location of part B. Background is the Ocean base map from ESRI et al. (2015). (B) Generalized geologic map of Lake Towuti and the surrounding area showing the dominance of ultramafic material. Map is modified from Costa et al. (2015). (C) Bathymetry of Lake Towuti showing the location of the two analyzed sediment cores at the distal margins of the Mahalona River delta (white dots labeled 4 for TOW4 and 5 for TOW5). River inputs are shown in thin blue lines, with the Mahalona River shown as a thick blue line. Credit: Goudge et al. and GSA Bulletin

In their *GSA Bulletin* article published online last week, Timothy A. Goudge and colleagues detail the clay mineralogy of sediment from Lake Towuti, Indonesia, using a technique called visible to near-infrared (VNIR) spectroscopy. VNIR measures the signature of reflected light from a sample across a larger wavelength range than just visible light. At Lake Towuti, the spectral record shows distinct variations in clay mineralogy over the past 40,000 years.

The record also captures the response of the lake system to changing climate, including changes in lake levels, forced delta progradation, and river incision. According to Goudge and colleagues, this demonstrates the utility of VNIR spectroscopy in developing paleoenvironmental records over tens of thousands of years.

Interestingly, Goudge and colleagues also suggest that paleolake deposits on Mars should preserve similar paleoenvironmental information that could be accessed through remote sensing studies of stratigraphy and VNIR reflectance spectroscopy.

Lead author Tim Goudge says, "The major link between this study of [lake sediment](#) in Indonesia and lake deposits on Mars is in terms of the methods we used. We can differentiate material properties with color (e.g., rust is red because of the iron in it), so VNIR spectroscopy allows us to determine what minerals are within a sample of lake sediment."

While this technique is relatively new for applying to lake deposits on Earth, it can be run remotely simply using reflected sunlight. Thus it is commonly employed to study the mineralogy and composition of the surface of other planetary bodies, including Mars.

Goudge says, "Our study shows that one can use VNIR [spectroscopy](#) to understand the evolution of past climate that is recorded by lake sediment. Therefore, we propose that applying a similar approach to studying ancient [lake](#) deposits on Mars at high resolution will help to unravel the history of ancient martian climate."

Lake Towuti is also contained within an ophiolite, which is composed of very iron- and magnesium-rich (mafic) rocks, and so is more comparable to the mafic surface of Mars than much of Earth's land surface, which is commonly more felsic (richer in silicon, aluminum, sodium, etc.). The compositional link is not perfect, however, so Goudge and colleagues more heavily emphasize the applicability of the technique.

**More information:** Timothy A. Goudge et al. A 40,000 yr record of clay mineralogy at Lake Towuti, Indonesia: Paleoclimate reconstruction from reflectance spectroscopy and perspectives on paleolakes on Mars, *Geological Society of America Bulletin* (2017). [DOI: 10.1130/b31569.1](https://doi.org/10.1130/b31569.1)

Provided by Geological Society of America

Citation: Paleolake deposits on Mars might look like sediments in Indonesia (2017, March 6)  
retrieved 3 May 2024 from

<https://phys.org/news/2017-03-paleolake-deposits-mars-sediments-indonesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.