

Satellites reveal bird habitat loss in California

March 27 2017



Satellite images have shown a dramatic loss of habitat for migratory shorebirds like these passing through California. Seasonally flooded farm fields could provide additional habitat for these birds. Credit: Danica Schaffer-Smith, Duke University

Drought and reduced seasonal flooding of wetlands and farm fields

threaten a globally important stopover site for tens of thousands of migratory shorebirds in California's Sacramento Valley, a new Duke University-led study shows.

The researchers' analysis of historical biweekly NASA Landsat satellite images of the valley reveals that flooded [habitat](#) near the peak time of spring migration has shrunk by more than twice the size of Washington, D.C. over the last 30 years.


"On average, we're losing an area about four times the size of Central Park each year, during a critical window of time in late March," said Danica Schaffer-Smith, a doctoral student at Duke's Nicholas School of the Environment, who conducted the study with researchers from the nonprofit Point Blue Conservation Science.

More than half of all shorebird species in the Western hemisphere are now in decline, Schaffer-Smith noted.

The Sacramento Valley site is an internationally important resting and refueling stop for tens of thousands of these wetland-dependent birds traveling along the Pacific Flyway. Some of these species migrate thousands of miles from Argentina to Alaska and back again each year.

"The fact that these highly mobile species are increasingly struggling to find flooded habitat on their migrations is an indicator that our freshwater wetland systems are in trouble," Schaffer-Smith said.

"Freshwater is essential for all life. Many other species rely on these same habitats, too."



This is Landsat satellite data showing the presence of water for spring months in the Sacramento Valley from 1983 to 2015. This frequency was calculated as the number of times water was identified in each pixel, corrected by the total number of cloud-free instances. Credit: *Remote Sensing of Environment*, May 2017, pgs. 180-192.

The Sacramento Valley once supported a huge network of wetlands connected to the San Francisco Bay Delta, but more than 90 percent of them have been drained for agriculture. Analysis of recent satellite images by Schaffer-Smith and her team shows that open water covers just three percent of the landscape during peak migration in April, when the birds urgently need flooded habitat to rest and feed.

This winter's heavy rains are unlikely to reverse these habitat losses.

"One season of plentiful rainfall can't undo the effects of years of habitat destruction and increased water consumption for a growing number of competing uses," she said. "However, most water in California is captured in reservoirs and diverted through a system of canals and ditches, which also makes it possible for us to intentionally manipulate flooded habitat on the landscape."

During the worst of the recent drought years, conservation organizations joined forces to launch BirdReturns, a payment-for-services program that compensated farmers for flooding their fields to provide additional habitat for birds, Schaffer-Smith said. The new study's findings could help guide the future timing and location of such initiatives.

"Years of drought have heightened scrutiny of water use in California to the point that even rice farmers have begun to explore a switch to drip irrigation to conserve water, but these fields provide important habitat where wetlands have been lost," she said.

Schaffer-Smith and her colleagues published their peer-reviewed paper this month in the journal *Remote Sensing of Environment*. The study is freely available online through May 3, 2017.

"Satellite imagery can help us get the biggest bang for our buck by targeting conservation initiatives in a specific window of time at key locations," she said. "Landsat is the longest running Earth observation satellite system we have, and free access to this data enables researchers to look at the effects of seasonality, climate cycles, and long-term trends in land-use change."

More information: Danica Schaffer-Smith et al, Three decades of Landsat-derived spring surface water dynamics in an agricultural wetland mosaic; Implications for migratory shorebirds, *Remote Sensing of Environment* (2017). [DOI: 10.1016/j.rse.2017.02.016](https://doi.org/10.1016/j.rse.2017.02.016)

Provided by Duke University

Citation: Satellites reveal bird habitat loss in California (2017, March 27) retrieved 25 April 2024 from <https://phys.org/news/2017-03-satellites-reveal-bird-habitat-loss.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.