


As sea level rises, much of Honolulu and Waikiki vulnerable to groundwater inundation

March 28 2017



A graphic showing how sea level rise lifts freshwater, causing groundwater inundation in low-lying areas. Credit: UHM Coastal Geology Group

New research from the University of Hawai'i at Mānoa (UHM) reveals a large part of the heavily urbanized area of Honolulu and Waikiki, Hawai'i is at risk of groundwater inundation—flooding that occurs as

groundwater is lifted above the ground surface due to sea level rise. Shellie Habel, lead author of the study and doctoral student in the UHM Department of Geology and Geophysics, School of Ocean and Earth Science and Technology (SOEST), and colleagues developed a computer model that combines ground elevation, groundwater location, monitoring data, estimates of tidal influence, and numerical groundwater-flow modeling to simulate future flood scenarios in the urban core as sea level rises three feet, as is projected for this century under certain climate change scenarios.

"This flooding will threaten \$5 billion of taxable real estate; flood nearly 30 miles of roadway; and impact pedestrians, commercial and recreation activities, tourism, transportation, and infrastructure," said Habel. "The flooding will occur regardless of seawall construction, and thus will require innovative planning and intensive engineering efforts to accommodate standing water in the streets."

Surprisingly, the team of researchers also discovered 86% of active cesspools in the study area are likely currently inundated by groundwater. This suggests that cesspool effluent is now entering coastal groundwater and coastal environments in the study area. Sea level rise of approximately three feet would fully inundate 39 cesspools, introducing effluent at the ground surface where people work and live. This presents a serious health concern that will become progressively more serious as contaminated waters begin breaching the ground surface.

They also found that the water table is close to the ground surface—within two feet at high tide—in many places. This narrow unsaturated space means that groundwater inundation will become a serious concern well before the end of the century. When it rains and infiltration fills this space, it is a problem already.

"Waikiki, the gateway of the state's tourism industry, currently has such


narrow unsaturated space that many construction projects working below the ground surface have to dewater the excavation before construction can begin," said Habel.

"Our findings suggest that coastal communities in Hawai'i and globally are exposed to complex [groundwater](#) flooding hazards associated with [sea level](#) rise in addition to the typical concerns of coastal erosion and wave overtopping," said Chip Fletcher, professor of Geology and Geophysics and associate dean in SOEST and principal investigator on the study. "Groundwater inundation will require entirely unique adaptation methods if we are to continue to live in and develop the coastal zone. Coastal planners and community stakeholders will need to work with architects, engineers, geologists, ecologists, economists, hydrologists, and other innovative thinkers in order to manage these problems."



This construction trench in Waikiki shows that the water table is nearly at the ground surface at high tide. Credit: UHM Coastal Geology Group

This study identified particular locations and infrastructure that will be vulnerable to future flooding and is a crucial first step towards addressing future challenges. The team of researchers hope to use this methodology to identify future flooding and at risk infrastructure in other locations, as well as assist in developing adaptation efforts among vulnerable coastal communities.



Simulations of groundwater inundation (blue) and narrow unsaturated space (yellow) today and considering sea level rise of approximately 1, 2 and 3 ft during an average spring (monthly maximum) tide stage. Credit: UHM Coastal Geology Group

More information: Shellie Habel et al, Development of a model to simulate groundwater inundation induced by sea-level rise and high tides in Honolulu, Hawaii, *Water Research* (2017). [DOI: 10.1016/j.watres.2017.02.035](https://doi.org/10.1016/j.watres.2017.02.035)

Provided by University of Hawaii at Manoa

Citation: As sea level rises, much of Honolulu and Waikiki vulnerable to groundwater inundation (2017, March 28) retrieved 2 February 2023 from <https://phys.org/news/2017-03-sea-honolulu-waikiki-vulnerable-groundwater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.