

Even short-duration heat waves could lead to failure of coffee crops

March 30 2017



Credit: Oregon State University

"Hot coffee" is not a good thing for java enthusiasts when it refers to plants beset by the high-temperature stress that this century is likely to bring, research at Oregon State University suggests.

A study by OSU's College of Forestry showed that when *Coffea arabica* plants were subjected to short-duration [heat](#) waves, they became unable to produce flowers and fruit.

That means no [coffee](#) beans, and no coffee to drink.

C. arabica is the globe's dominant coffee-plant species, accounting for 65 percent of the commercial production of the nearly 20 billion pounds of coffee consumed globally each year.

Continually producing new flushes of leaves year-round, *C. arabica* grows on 80 countries in four continents in the tropics.

The OSU research investigated how leaf age and heat duration affected *C. arabica*'s recovery from heat stress during greenhouse testing. A major finding was that the younger, "expanding" leaves were particularly slow to recover compared to mature leaves, and that none of the plants that endured the simulated heat waves produced any flowers or fruit.

"This emphasizes how sensitive *Coffea arabica* is to temperature," said lead author Danielle Marias, a plant physiologist with OSU's Department of Forest Ecosystems and Society. "No flowering means no reproduction which means no beans, and that could be devastating for a coffee farmer facing crop failure.

"Heat is very stressful to the plants and is often associated with drought. However, in regions where coffee is grown, it may not just be hotter and drier, it could be hotter and wetter, so in this research we wanted to isolate the effects of heat."

In the OSU study, *C. arabica* [plants](#) were exposed to heat that produced leaf temperatures of a little over 120 degrees Fahrenheit, for either 45 or 90 minutes. That leaf temperature, Marias emphasizes, is a realistic result of global climate change and also more than the surrounding air temperature – think of how hot, for example, asphalt gets in the sunshine on a 90-degree day.

Expanding leaves subjected to the 90-minute treatment took the longest to recover physiologically as measured by photosynthesis; chlorophyll

fluorescence, an indicator of photosynthetic energy conversion; and the presence of nonstructural carbohydrates, which include starch and free sugars involved in growth, reproduction and other functions.

"In both treatments, photosynthesis of expanding leaves recovered more slowly than in mature leaves, and stomatal conductance of expanding leaves was reduced in both heat treatments," Marias said. "Based on the leaf energy balance model, the inhibited stomatal conductance reduces evaporative cooling of leaves, which could further increase leaf temperatures, exacerbating the aftereffects of heat stress under both full and partial sunlight conditions, where *C. arabica* is often grown."

Regardless of [leaf](#) age, the longer heat treatment resulted in decreased water-use efficiency, which could also worsen the effects of [heat stress](#), particularly during drought.

More information: Danielle E. Marias et al. Impacts of leaf age and heat stress duration on photosynthetic gas exchange and foliar nonstructural carbohydrates in, *Ecology and Evolution* (2017). [DOI: 10.1002/ece3.2681](#)

Provided by Oregon State University

Citation: Even short-duration heat waves could lead to failure of coffee crops (2017, March 30) retrieved 3 May 2024 from <https://phys.org/news/2017-03-short-duration-failure-coffee-crops.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|