

# Subaru telescope detects shadow of gas cloud in ancient proto-supercluster

March 28 2017



The distribution of galaxies in the proto-supercluster region 11.5 billion years ago (top left), and the Subaru Telescope Suprime-Cam image used in this work (right, larger image). Neutral hydrogen gas distribution is superposed on the Subaru image. The red color indicates denser regions of the neutral hydrogen gas. Cyan squares correspond to member galaxies in the proto-supercluster, while objects without cyan squares are foreground galaxies and stars. The



distribution of neutral hydrogen gas does not align perfectly with the galaxies. Credit: Osaka Sangyo University / NAOJ

A team led by researchers from Osaka Sangyo University, with members from Tohoku University, Japan Aerospace Exploration Agency (JAXA) and others, has used the Suprime-Cam on the Subaru Telescope to create the most-extensive map of neutral hydrogen gas in the early universe. This cloud appears widely spread out across 160 million light-years in and around a structure called the proto-supercluster. It is the largest structure in the distant universe, and existed some 11.5 billion years ago. Such a huge gas cloud is extremely valuable for studying large-scale structure formation and the evolution of galaxies from gas in the early universe, and merits further investigation.

"We are surprised because the dense gas <u>structure</u> is extended much more than expected in the proto-supercluster," said Dr. Mawatari. "Wider field observations with narrow-band filters are needed to grasp full picture of this largest structure in the young Universe. This is exactly the type of strong research that can be done with Hyper Suprime-Cam (HSC) recently mounted at the Subaru Telescope. We intend to study the gas - galaxy relation in various proto-superclusters using the HSC."

## **Understanding Matter Distribution in the Universe**

Stars assembled to form galaxies, and galaxies are clustered to form larger structures such as clusters or superclusters. Matter in the current universe is structured in a hierarchical manner on scales of ~ 100 million light-years. However, we cannot observe inhomogeneous structure in any direction or distance over scales larger than that. One important issue in modern astronomy is to clarify how perfectly the large-scale uniformity and homogeneity in matter distribution is maintained. In addition,



astronomers seek to investigate the properties of the seeds of large-scale structures (i.e., the initial matter fluctuations) that existed at the beginning of the universe. Thus, it is important to observe huge structures at various epochs (which translates to distances). The study of gaseous matter as well as galaxies is needed for an accurate and comprehensive understanding. This is because local superclusters are known to be rich in gas. Furthermore, it is clear that there are many newborn galaxies in ancient (or distant) clusters. A detailed comparison between the spatial distributions of galaxies and gas during the early epochs of the universe is very important to understand process of galaxy formation from the dim (low light-emitting) clumps of gas in the early universe.



Schematic pictures of an analysis scheme of previous work (left) and a new method (right). In the previous approach, basically a single background light source (quasar) can be used in a searched area. On the other hand, with the new scheme, it is easier to spatially resolve the neutral hydrogen gas density by using many normal galaxies in a searched area as background light sources. In the new



scheme, absorption strength by the neutral hydrogen gas is estimated by measuring how much flux of the background galaxies becomes dimmed in the narrow-band image, not by using spectrum. By combining this scheme with the wide-area imaging ability of the Subaru Telescope, Mawatari, et al. made the most-extensive map of neutral hydrogen gas ever created. Credit: Credit: Osaka Sangyo University / NAOJ

In order to investigate early, dim gas clouds, astronomers take advantage of the fact that light from bright distant objects gets dimmed by foreground gas (giving an effect like a "shadow picture"). Since neutral hydrogen in the gas cloud absorbs and dims light from background objects at a certain wavelength, we can see characteristic absorption feature in the spectrum of the background object. In many previous observations, researchers used quasars (which are very bright and distant) as background light sources. Because bright quasars are very rare, opportunities for such observations are limited. This allows astronomers to get information about the gas that lies only along the line of sight between a single QSO and Earth in a wide survey area. It has long been the goal to obtain "multi-dimensional" information of gas (e.g., spatially resolve the gas clouds) rather than the "one-dimensional" view currently available. This requires a new approach.

## **Expanding the View**

To widen their view of these objects in the early universe, Dr. Ken Mawatari at Osaka Sangyo University and his colleagues recently developed a scheme to analyze the spatial distribution of the <u>neutral</u> <u>hydrogen gas</u> using imaging data of galaxies of the distant epoch. There are two major advantages to this approach. First, instead of rare quasars, the team uses numerous normal galaxies as background light sources to investigate gas distribution at various places in the search area. Second,



they use imaging data taken with the narrow-band filter on Suprimecam. It is fine-tuned so that light with certain wavelengths can be transmitted, to capture evidence of absorption by the neutral hydrogen gas (the shadow picture effect). Compared with the traditional scheme of observations based on spectroscopy of quasars, this new method enables Mawatari and his collaborators to obtain wide-area gas distribution information relatively quickly.

The researchers applied their scheme to the Subaru Telescope Suprime-Cam imaging data taken in their previous large survey of galaxies. The fields investigated in this work include the SSA22 field, an ancestor of a supercluster of galaxies (proto-supercluster), where young galaxies are formed actively, in the universe 11.5 billion years ago in the early universe.





Sky distribution of the neutral hydrogen gas in the three fields studied in this work. While in the normal fields (SXDS and GOODS-N) the neutral hydrogen gas density is consistent with the average density in the entire universe at 11.5 billion years ago, the neutral hydrogen gas density is higher than the average over the entire SSA22 proto-supercluster field. Contours correspond to the galaxies $\hat{a} \in \mathbf{O}$  number density. Bold, solid thin, and dashed contours mean the average, high density, and low density regions, respectively. Credit: Osaka Sangyo University / NAOJ

#### New Maps of Neutral Hydrogen Distribution

The researchers' work resulted in very wide-area maps of the neutral hydrogen gas in the three fields studied. It appears that the neutral hydrogen gas absorption is significantly strong over the entire SSA22 proto-supercluster field compared with those in the normal fields (SXDS and GOODS-N). It is clearly confirmed that the proto-supercluster environment is rich in neutral hydrogen gas, which is the major building block of galaxies.

The team's work also revealed that gas distribution in the protosupercluster region does not align with the galaxies' distribution perfectly. While the proto-supercluster is rich in both galaxies and gas, there is no local-scale dependency of gas amount correlated with the density of galaxies inside the proto-supercluster. This result may mean that the neutral hydrogen gas not only is associated with the individual galaxies but also spreads out diffusely across intergalactic space only within the proto-supercluster. Since the neutral hydrogen gas excess in the SSA22 field is detected over the entire searched area, this overdense gas structure is actually extended more than 160 million light-years. In the traditional view of structure formation, matter density fluctuation is



thought to be smaller and large-scale high-density structure was rarer in the <u>early universe</u>. The discovery that a gas structure that extends across more than 160 million light-years (which is roughly same as present-day superclusters in scale) already existed in the universe 11.5 billion years ago is a surprising result of this study.

By investigating spatial distribution of the neutral hydrogen gas in a very large area, the researchers have provided a new window on the relation between gas and <u>galaxies</u> in the young <u>universe</u>. The SSA22 huge gas structure revealed by this work is considered a key object to test the standard theory of structure formation, and so further investigation is anticipated.

**More information:** Ken Mawatari et al, Imaging of diffuse H i absorption structure in the SSA22 proto-cluster region at z = 3.1, *Monthly Notices of the Royal Astronomical Society* (2017). DOI: <u>10.1093/mnras/stx038</u>

#### Provided by National Institutes of Natural Sciences

Citation: Subaru telescope detects shadow of gas cloud in ancient proto-supercluster (2017, March 28) retrieved 3 May 2024 from <u>https://phys.org/news/2017-03-subaru-telescope-shadow-gas-cloud.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.