

How vision may have driven fishes onto land

March 8 2017, by John Long



375 million years ago fishes like Tiktaalik (pictured, above) looked out above water for prey. Credit: John Long, Flinders University, Author provided

About 375 million years ago, certain fishes had developed powerfully strong paired fins that were capable of transporting them out of the water and onto land.

These fishes would eventually evolve into the first truly terrestrial animals, called <u>tetrapods</u>. They had four limbs bearing digits – fingers and toes – to help them when they walked around around on land.



But one of the biggest mysteries for scientists is figuring out what could have driven such fishes out of the water and onto land in the first place.

Was it availability of new food sources, or perhaps their need to escape from predators in the water?

A new theory says it was improved vision, as shown by dramatic increases in eye size and visual acuity, that enabled fishes peeping upwards at the waterline to spot prey on land.

This would have motivated them to venture out of the water to hunt for food. The new research is <u>published today</u> by <u>Malcolm MacIver</u> from Northwestern University in the US and colleagues. They have named this the "Buena Vista" hypothesis, from the Spanish for "good view".

All the better to see you with

The team measured the eye sizes of many kinds of <u>fossil fishes</u> and <u>early</u> <u>tetrapods</u> that lived between 390 million and 250 million years ago to show a dramatic increase in eye size just before the most advanced fishes, called elpistostegalians, left the water to evolve into the first tetrapods.

While fossil eyeball capsules are sometimes preserved perfectly in 3-D, most fossil eyes are reconstructed from the shape of the curved <u>sclerotic</u> ring bones that surround the eye, combined with size and shape of the orbit, which is the hole in the skull that envelops the eyeball.

The increase in eye-socket size as measured from fossil skulls of fishes and early amphibians corresponds to a three-fold increase in eye size.

Experimental models of visual ability against different environment scenarios, from deep in water to being in air during daylight, give an



estimation of the efficiency of vision in each case. The researchers computed the range, volume and viewing factors across a range of conditions and pupil sizes.

The results showed that fishes underwent a series of visionary improvements in stages moving from water to land. The first step was when they began to look upwards towards light. Ambush predators, which surprise their prey by a rapid attack coming up from underneath, do this.

A Devonian <u>fish</u> called *Panderichthys* had a head shaped perfectly for this type of attack. Its vision was best adapted for scanning across the plane of view when looking up for prey.

The second stage was when fishes took to surfacing and poking their eyes above water to peer through the air at the environment around them.

Fishes such as *Tiktaalik*, the fish most closely related to tetrapods, exemplify this stage very well. It has a long crocodile like head with similarly placed eyes in the centre, raised up so they can look around.

It was found that after the emergence of vision, from the fully aquatic vision to totally aerial vision, there was a five million fold increase in the total amount of space monitored by vision.





How the eye size (white circle) increased dramatically in the evolution of lobefinned fish that lived entirely in water to the first land living tetrapods. Credit: John Long, Flinders University

Even looking from the water's edge, by poking their eyes above the water, fishes such as *Tiktaalik* had a massive increase in visual ability to spot prey, such as large worms or bugs on the nearby river bank.

The latest work is not without some doubts, as lead author MacIver told me:

The simulations showed that the bulk of the eye size increase occurred before tetrapods were on land, while my own earlier prediction was that the increase would occur after they were on land.

That's because I'd only envisaged seeing through air happening at the same time as terrestriality. So it was a big relief when I subsequently came across Long and Gordon's 2004 paper that synthesized several anatomical changes and gave the idea that early tetrapods, still largely fully aquatic, may have hunted like crocodiles – living in water but



hunting by seeing through air.

The new research suggests that this Buena Vista view was perhaps the zip-line effect for fishes to take more active forays onto land. This would have accelerated the development of more robust limbs and the emergence of well-developed digits to replace fin rays.

Some of these advanced fishes, such as *Panderichthys*, <u>already had digits</u> present inside their fins.

Fully formed rows of fingers and toes would be clearly more advantageous for moving on land, though some of the first tetrapods had evolved digits whilst living in water.





How vertebrate vision improved dramatically in the stages from life in water as a fish to life on land as a limbed tetrapod. Credit: Malcom MacIver, Northwestern University

Vision and the beginning of complex planning

The big implication of this research is the effect it all had on our brains with the shift in behaviour and cognition that accompanied this transition onto land.



With larger eyes, vertebrates began long range sensing for the first time. To sneak up on prey successfully implies they must have been planning an ambush attack, such as creeping up from behind. Otherwise the prey would see them coming and escape if they approached directly.

In living fishes and amphibians the time between prey detection and escape is reduced by about 6 microseconds through using a single large cell called a Mauthner neuron cell to initiate the escape plan. This neural circuity enables ultra-fast reactions to stimuli.

These Mauthner command cells are today only found in fishes and amphibians. Later, more efficient nerves in the spine displaced the need for Mauthner cells.

It seems that the origin of long range vision started with a common ancestor of fishes and <u>tetrapods</u>, most likely one of the highly advanced lobe-finned fishes such as *Tiktaalik*.

So be thankful for our fishy ancestors that boldly ventured to go where no fish had gone before, out of the <u>water</u> up on to land. Without the long range vision that accompanied this event our ability to plan and perform complex actions might never have evolved.

This article was originally published on <u>The Conversation</u>. Read the <u>original article</u>.

Provided by The Conversation

Citation: How vision may have driven fishes onto land (2017, March 8) retrieved 16 July 2024



from https://phys.org/news/2017-03-vision-driven-fishes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.