

Using an airplane to measure volatile organic compounds from oil sands surface mining facilities

April 25 2017, by Bob Yirka



Aerial view of Fort McMurray. Credit: Regional Municipality of Wood Buffalo/ Wikipedia

(Phys.org)—A team of researchers affiliated with several institutions in

Canada has found that using an airplane to carry emissions reading hardware over mining facilities offers a better way to measure volatile organic compounds (VOCs) being pumped into the atmosphere than methods currently in use. In their paper published in *Proceedings of the National Academy of Sciences*, the team describes what they found when they used their flying emissions reading lab over a real oil sands surface mining facility and why they believe similar approaches should be used for other facilities around the world.

As difficult as it may be to believe, governmental agencies in the U.S. and Canada both rely on self-reporting by oil companies to determine if such companies are complying with federal regulations when making policy regarding reversing the impact of global warming. As the researchers note, such data is typically obtained by installing monitoring devices on smokestacks or other obvious sources of emissions—but they do not measure emissions from leaks and other operational areas—instead they use math, extrapolating data from other substance emissions to make estimates of VOC emissions. But as the researchers also note, to date, the means by which oil companies come up with their data has not been tested. In this new effort, the research team sought to do just that by installing air monitoring equipment on an airplane and then flying over a mining facility and taking air samples, testing them, and then comparing what they found with statistics given by the [oil company](#).

To get good readings, the researchers flew in a 275-square kilometer box pattern at different altitudes over a surface mining facility collecting samples every hour for several hours. The samples were then returned to their lab for analysis. The team reports that their readings were from two to 4.5 times higher than the readings given by the [oil company](#), suggesting that the company was vastly underreporting the amount of VOCs it emits.

The researchers note that their technique could be used on a wide variety of [mining](#) operations, including hydraulic fracturing and oilsands projects. It could also be modified to test for more than just VOCs to get a better picture of all the pollutant types emitted at a given site.

More information: Shao-Meng Li et al. Differences between measured and reported volatile organic compound emissions from oil sands facilities in Alberta, Canada, *Proceedings of the National Academy of Sciences* (2017). [DOI: 10.1073/pnas.1617862114](https://doi.org/10.1073/pnas.1617862114)

Abstract

Large-scale oil production from oil sands deposits in Alberta, Canada has raised concerns about environmental impacts, such as the magnitude of air pollution emissions. This paper reports compound emission rates (E) for 69–89 nonbiogenic volatile organic compounds (VOCs) for each of four surface mining facilities, determined with a top-down approach using aircraft measurements in the summer of 2013. The aggregate emission rate (aE) of the nonbiogenic VOCs ranged from 50 ± 14 to 70 ± 22 t/d depending on the facility. In comparison, equivalent VOC emission rates reported to the Canadian National Pollutant Release Inventory (NPRI) using accepted estimation methods were lower than the aE values by factors of 2.0 ± 0.6 , 3.1 ± 1.1 , 4.5 ± 1.5 , and 4.1 ± 1.6 for the four facilities, indicating underestimation in the reported VOC emissions. For 11 of the combined 93 VOC species reported by all four facilities, the reported emission rate and E were similar; but for the other 82 species, the reported emission rate was lower than E. The median ratio of E to that reported for all species by a facility ranged from 4.5 to 375 depending on the facility. Moreover, between 9 and 53 VOCs, for which there are existing reporting requirements to the NPRI, were not included in the facility emission reports. The comparisons between the emission reports and measurement-based emission rates indicate that improvements to VOC emission estimation methods would enhance the accuracy and completeness of emission estimates and their applicability

to environmental impact assessments of oil sands developments.

© 2017 Phys.org

Citation: Using an airplane to measure volatile organic compounds from oil sands surface mining facilities (2017, April 25) retrieved 23 April 2024 from <https://phys.org/news/2017-04-airplane-volatile-compounds-oil-sands.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.