

# How domestication can change animals' facial features

April 10 2017, by Antje Karbe

---

Domesticated animals, compared to their wild counterparts, have undergone numerous changes in physiology, behavior and morphology. These changes are commonly referred to as the domestication syndrome and include behavioral changes, such as increased docility as well as genetic alterations in size, color and facial characteristics. In attempting to find whether these changes have a single cause, Russian zoologist Dmitry Belyaev conducted a series of selection experiments with silver foxes, hypothesizing that behavior, specifically tameness, was the key driving factor behind the changes brought about by domestication. After generations of selecting foxes for tameness, they were found to display phenotypes similar to those observed in domesticated species. Since then, it has been further hypothesized that selection for social tolerance and reduced aggression may also have played an important role in shaping the modern human anatomy, which is remarkable for the reduced face and gracile overall features.

In parallel to his fox experiment, Belyaev also selected rats over 64 generations for their behavior: either tameness or defensive aggression towards humans. In the first ever quantitative study on the facial anatomy of Belyaev's selected rats, an international team of researchers from the Senckenberg Center for Human Evolution and Paleoenvironment at the University of Tübingen and Pennsylvania State University collected 3-D measurements on the skulls of both tame- and aggressive-selected rats, in order to evaluate Belyaev's hypothesis that tame behavior correlates with the facial changes similar to those seen in domesticated animals. The study found that rats selected for tame

behavior show some – though not all- traits present in domesticated animals and the tame silver foxes. The findings are published in the latest PlosOne.

Senior author, Tübinger palaeoanthropologist Professor Katerina Harvati and Dr. Nandini Singh from Pennsylvania State University conceived the project, in collaboration with Dr. Frank Albert from the University of Minnesota. Quantitative analyses of cranial measurements revealed clear differences, including a smaller and retracted snout, but no difference in overall size, between rats selected for either tameness or defensive aggression. However, unlike Belyaev who found a kind of "feminization" and sexual dimorphism in the tame silver foxes – traits also seen in domesticated [animals](#) - Harvati and colleagues did not see this trend among the tame [rats](#).

Professor Harvati says this shows that not all elements of the domestication syndrome might appear in [domesticated animals](#), depending on the species. However, this study con-firms the wide-ranging effect of [behavioral changes](#) on the phenotype over generations. Experimental animal models can provide novel ways to test and address a number of questions regarding the developmental origins of animal domestication not otherwise possible with archaeological samples.

**More information:** Nandini Singh et al. Facial shape differences between rats selected for tame and aggressive behaviors, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0175043](https://doi.org/10.1371/journal.pone.0175043)

Provided by University of Tübingen

Citation: How domestication can change animals' facial features (2017, April 10) retrieved 30 April 2024 from <https://phys.org/news/2017-04-domestication-animals-facial-features.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.