


Feeding fat to fungi: Evidence for lipid transfer in arbuscular mycorrhiza

April 6 2017



Confocal microscopy image of fungal arbuscules (green) inside plant root cells (red). Credit: Dr. Maria J. Harrison, BTI

Nearly all organisms live in symbiosis with a vast, diverse array of microbes. Arbuscular mycorrhizal (AM) symbiosis is the interaction between plants and a group of fungi called Glomeromycota. Most land plants, including several crop species, are able to interact with these fungi, which have been long known to positively affect plant growth and nutrition. The fungi live in plant roots where they elongate their tendrils

(called hypha) into the surrounding soil, like an extension of the root system, to better access and transfer nutrients to the plant. In return the plant serves the fungus food made during photosynthesis.

Previous research has shown that an AM fungal dinner at least consists of [high-energy sugars](#), but published today in *New Phytologist*, researchers from the labs of Dr. Maria Harrison at the Boyce Thompson Institute and Dr. Peter Dörmann at the University of Bonn have produced the first experimental evidence to suggest that [AM fungi also get their fat \(lipids\) from the host plant](#).

"Until recently, it has been assumed that the [fungus](#) obtains sugar from the plant and can manage to make other [essential nutrients](#) itself," Harrison explains. "However, this is not the full story."

The transfer of [lipid](#) may be just as important. Living inside a plant root, the fungus will enter the root cells and create a tree-like structure called an arbuscule, giving the [symbiosis](#) its namesake. As an arbuscule is formed, new plant and fungal membrane must be generated, like the bark on the extending branches of a growing tree. To do so both organisms ramp up production of lipids, which are important building blocks for cell membranes. Sure enough, plant root cells that harbor fungal arbuscules [increase their expression of genes involved in producing such lipids](#) by 3000-fold.

Taking it a step further, Bravo et al. showed that individually mutating two of those genes, *FatM* and *RAM2*, in the model legume *Medicago truncatula* not only disrupts the fungus' ability to form healthy arbuscules, but also reduces plant production of a lipid necessary for fungal growth, called 16:0 β -monoacylglycerol (MAG). This result indicates that *FatM* and *RAM2* may play specific roles in the biosynthesis of 16:0 β MAG, and further highlights the importance of this particular lipid for AM symbiosis.

What makes this food exchange even more important is the fact that AM fungi can't synthesize 16:0 β MAG themselves, indicating the [fungi](#) depend on their [host plant](#) as the source of this molecule. The exciting discovery that the plant provides the fungus with this essential lipid helps explain the fungus' mysterious obligate nature, meaning it is completely dependent on its host for survival and reproduction.

"Without the basic lipids that are obtained from the plant, the AM fungus cannot produce the complex lipids that it requires to live," says first author Armando Bravo.

One of the next steps in this research is to investigate how the plant regulates 16:0 β MAG production, and whether more fat for the fungus means more nutrient transfer to the plant in return. In a world where agricultural lands are being depleted of essential nutrients, AM symbiosis may be the key to healthier [plants](#) and greater crop yield.

More information: Armando Bravo et al, Arbuscular mycorrhiza-specific enzymes FatM and RAM2 fine-tune lipid biosynthesis to promote development of arbuscular mycorrhiza, *New Phytologist* (2017). [DOI: 10.1111/nph.14533](https://doi.org/10.1111/nph.14533)

Provided by Boyce Thompson Institute

Citation: Feeding fat to fungi: Evidence for lipid transfer in arbuscular mycorrhiza (2017, April 6) retrieved 9 April 2024 from <https://phys.org/news/2017-04-fat-fungi-evidence-lipid-arbuscular.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
