

Study discovers fundamental unit of cell size in bacteria

April 13 2017



Biologists have long known that bacteria grow faster and bigger when the quality of nutrients becomes better, a principle in microbial physiology known as the "growth law," which describes the relationship between the average cell size of bacteria and how fast they grow.

But the growth law has a major hole: It is unable to explain why [bacteria](#)

divide when they reach a certain critical size, no matter how much or how little nutrients are available.

By applying mathematical models to a large number of experiments in which bacterial growth is inhibited, however, a team of physicists, biologists and bioengineers from UC San Diego discovered the reason for this and in the process developed a "general growth law" that explains the origin of these idiosyncrasies of bacterial physiology.

The researchers detailed their achievements in a paper published in this week's issue of the journal *Current Biology*.

"A few years ago, we set out to do extensive growth inhibition experiments to test the growth law using the model organism *Escherichia coli*," said Suckjoon Jun, an assistant professor of physics and molecular biology at UC San Diego, who headed the research effort. "Perhaps not so surprisingly, the original growth law was unable to explain changes in [cell size](#) under growth inhibition. Cell size either increased or decreased depending on the inhibition method. Sometimes, cell size did not change at all despite significant growth inhibition."

Jun and his colleagues discovered that when cells began replicating their genetic material in preparation for cell division, cell size remained remarkably constant despite the many genetic processes and changes in the cell such as protein and DNA synthesis, [cell wall synthesis](#) and cell shape.

"We realized that this invariant cell size represents a fundamental unit of cellular resources required to start growth and the cell cycle, or the 'engine' of a car, so to speak," said Jun. "This 'unit cell' is the fundamental building block of cell size, and cell size is the sum of all invariant unit cells for any growth condition, explaining the origin of the growth law."

Jun said the development of high-throughput cell sampling techniques and genetic methods such as "CRISPR interference" made it possible for his team to extract large amounts of physiological data from 10 million bacterial [cells](#) in their growth inhibition experiments.

"This allowed detailed and reliable statistics, and led to quantitative modeling that made experimentally testable predictions, helping us to understand the data at a deeper level," he added. "This complements the unexpected 'adder' principle that we discovered a few years ago."

Jun said this process was similar to the manner in which the Danish astronomer Tycho Brahe, by collecting better data of planetary orbits, was able to convince the German astronomer Johannes Kepler four centuries ago that planetary orbits, whose origin is gravity, were ellipses and not circles.

"Kepler's elliptical model said nothing about the physical origins of ellipses, but his kinematic modeling was an essential starting point for Newton's work on dynamics 50 years later," Jun said. "We don't know whether biology is following the footsteps of the history of physics, but examples are accumulating that some branches of biology are becoming as quantitative a science as physics."

Provided by University of California - San Diego

Citation: Study discovers fundamental unit of cell size in bacteria (2017, April 13) retrieved 13 March 2024 from <https://phys.org/news/2017-04-fundamental-cell-size-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
