


GERDA experiment ready to discover rarest radioactive decay

April 6 2017



View from bottom into the GERDA experiment: The fiber shroud of the liquid argon veto and the copper. Credit: V. Wagner/GERDA collaboration

Why is there more matter than antimatter in the universe? The reason might be hidden in the neutrino nature: one of the preferred theoretical

models assumes, that these elementary particles were identical with their own anti-particles. This in turn would lead to an extremely rare nuclear decay process, the neutrinoless double-beta decay ($0\nu\beta\beta$). The experiment GERDA now has reached a most important improvement in the search for $0\nu\beta\beta$ decay by reducing the disturbances (background) to an unprecedented low level making it the first "background-free" experiment in the field. This achievement is reported in the recent *Nature* article appearing April 6th, 2017.

Neutrinos are ghostly particles which are extremely hard to detect. They play a central role in how the sun burns, how supernovae explode and how elements are formed during the big bang. Determining their properties has advanced our understanding of [elementary particles](#) considerably, best documented by the fact that so far four Nobel prizes have been awarded to neutrino related research. One fundamental property is still unknown: are [neutrinos](#) Majorana particles, i.e. identical to their own anti-particles? In that case $0\nu\beta\beta$ decay will exist. Strong theoretical arguments favor this possibility and the above mentioned absence of anti-matter in our universe is likely connected to the Majorana character of neutrinos.

"Normal" [double beta decay](#) is an allowed rare process where two neutrons in a nucleus decay simultaneously into two protons, two electrons and two anti-neutrinos. It has been observed for some nuclei like ^{76}Ge , where single beta decay is not possible. The electrons and anti-neutrinos leave the nucleus, only the electrons can be detected. In $0\nu\beta\beta$ decay, no neutrinos leave the nucleus and the sum of the energies of the electrons is identical to the well known energy release of the decay. Measurement of exactly this energy is the prime signature for $0\nu\beta\beta$ decay.



Preparation of the GERDA experiment: Lowering the germanium detector array into the liquid argon tank - view from top.

Because of the importance of $0\nu\beta\beta$ decay in revealing the character of neutrinos and new physics, there are about a dozen experiments worldwide using different techniques and isotopes. The GERDA experiment is one of the leading experiments in the field, conducted by a European Collaboration. It is located in the underground Laboratori Nazionali del Gran Sasso of the Italian research organization INFN.

GERDA uses high-purity germanium detectors enriched in the isotope ^{76}Ge . Since the germanium is source and detector at the same time, a compact setup with minimum additional materials can be realized leading to low backgrounds and high detection efficiency. The excellent

energy resolution of germanium detectors and the novel experimental techniques developed by the GERDA collaboration provide unprecedented suppression of disturbing events from other radioactive decays (background events). Since $0\nu\beta\beta$ decay has a half-life many orders of magnitude longer than the age of the universe, the reduction of background events is most crucial for the sensitivity.

The bare germanium detectors are operated in 64 m^3 of liquid argon at a temperature of -190 degree Celsius. The argon container itself is inside a 590 m^3 tank filled with pure water which in turn is shielded by the Gran Sasso mountain against cosmic rays. The used argon and water are extremely pure in uranium and thorium; the liquids act as further shield for natural radioactivity from the surrounding. Their instrumentation provides additional means of background identification.

The novel techniques employed by GERDA reduced the number of background events in such a way, that now it is the first "background-free" experiment in the field. No $0\nu\beta\beta$ decays have been observed during the first five months of data taking and a lower half-life limit of 5×10^{25} yr was derived. Until the end of data taking in 2019 no background event should be left in the energy region where the $0\nu\beta\beta$ signal is expected and a sensitivity of 10^{26} yr will be reached. This makes GERDA best suited to discover a signal, which would manifest itself by a small number of events at the signal energy.

More information: M. Agostini et al. Background-free search for neutrinoless double- β decay of 76Ge with GERDA, *Nature* (2017). [DOI: 10.1038/nature21717](https://doi.org/10.1038/nature21717)

Data and Pictures: www.mpi-hd.mpg.de/gerda/public/index.html

Provided by Max Planck Society

Citation: GERDA experiment ready to discover rarest radioactive decay (2017, April 6) retrieved 20 April 2024 from <https://phys.org/news/2017-04-gerda-ready-rarest-radioactive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.