

Global warming accounts for tripling of extreme West African Sahel storms, study shows

April 26 2017



The onset of a Sahelian storm. Credit: Françoise GUICHARD/Laurent KERGOAT/CNRS

The UK-based Centre for Ecology & Hydrology (CEH) has led an international team of scientists who reveal global warming is responsible for a tripling in the frequency of extreme West African Sahel storms observed in just the last 35 years.

Professor Christopher Taylor, a Meteorologist at CEH, and researchers from partner institutions including Université Grenoble Alpes in France, also suggest that climate change will see the Sahel experience many more instances of extreme rain in future.

Professor Taylor and the fellow scientists' findings—published in the journal *Nature*—note that further strengthening of intense storms in the Sahel known as Mesoscale Convective Systems (MCSs) will increase the risk of more frequent and severe flooding and disease due to poor sanitation in West African cities. The findings will also this week be presented at the General Assembly of the European Geosciences Union at its meeting in Vienna, Austria.

The Sahelian storms are some of the most explosive storms in the world, containing clouds that can grow to a height of 16km above the ground. In 2009 a downpour of 263mm over several hours forced 150,000 residents of Ouagadougou, in Burkina Faso, to leave their homes. The study, which has analysed trends from 35 years of satellite observations across Africa, provides unique insight into how some of the most violent storms in the world are responding to rising global temperatures.



The onset of a Sahelian storm. Credit: Françoise GUICHARD/Laurent KERGOAT/CNRS

The research indicates that MCS intensification is linked to increasingly hot conditions in the Sahara desert resulting from man-made [greenhouse gas emissions](#). Saharan warming affects [storm](#) intensity across the Sahel, a band of semi-arid land to the south of the desert which is home to some of the most vulnerable populations on the planet.

Professor Taylor, said, "Global warming is expected to produce more intense storms, but we were shocked to see the speed of the changes taking place in this region of Africa."

Co-author Professor Douglas Parker, Professor of Meteorology at the University of Leeds, UK, said, "African storms are highly organised meteorological engines, whose currents extract water from the air to produce torrential rain. We have seen these engines becoming more efficient over recent decades, with resulting increases in the frequency of hazardous events."

Case study - flooding in Burkina Faso

Dr Abdoulaye Diarra, Senior Researcher at the International Institute for Water and Environmental Engineering in Ouagadougou, Burkina Faso:

"A total of 77 flood events have been recorded in Burkina Faso during the 31 year period from 1986 to 2016. The floods dynamic has increased in the Burkina from a frequency of 11 major events over 10 years (1.1 flood event/year) between 1986 to 2005 to a frequency of 55 over 11 years (5 flood events/year) from 2006 to 2016.

"The damages are generally more important in the urban area where runoff is very important and watercourses are limited. The 2009 flood (1 September), focused in Ouagadougou, caused heavy damage in the capital. More than 50 percent of the city's territory, including the main hospital of the city, was flooded.

"In total eight people died, more than 250 houses and 670 classrooms were destroyed, the main water purification plants for the city were out of use and nearly 150,000 people were affected within and around Ouagadougou."

More information: Christopher M. Taylor et al, Frequency of extreme Sahelian storms tripled since 1982 in satellite observations, *Nature* (2017). [DOI: 10.1038/nature22069](https://doi.org/10.1038/nature22069)

Provided by Centre for Ecology & Hydrology

Citation: Global warming accounts for tripling of extreme West African Sahel storms, study shows (2017, April 26) retrieved 10 April 2024 from <https://phys.org/news/2017-04-global-accounts-tripling-extreme-west.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--