


Quantum dots that emit infrared light open new window for biological imaging

April 10 2017



Researchers have found a way to make tiny particles that can be injected into the body, where they emit short-wave infrared light. The advance may open up a new way of making detailed images of internal body structures such as fine networks of blood vessels. Credit: Bawendi Group at MIT

For certain frequencies of short-wave infrared light, most biological

tissues are nearly as transparent as glass. Now, researchers have made tiny particles that can be injected into the body, where they emit those penetrating frequencies. The advance may provide a new way of making detailed images of internal body structures such as fine networks of blood vessels.

The new findings, based on the use of light-emitting particles called quantum dots, is described in a paper in the journal *Nature Biomedical Engineering*, by MIT research scientist Oliver Bruns, recent graduate Thomas Bischof PhD '15, professor of chemistry Mounqi Bawendi, and 21 others.

Near-infrared imaging for research on [biological tissues](#), with wavelengths between 700 and 900 nanometers (billionths of a meter), is widely used, but wavelengths of around 1,000 to 2,000 nanometers have the potential to provide even better results, because body tissues are more transparent to that light. "We knew that this imaging mode would be better" than existing methods, Bruns explains, "but we were lacking high-quality emitters"—that is, light-emitting materials that could produce these precise wavelengths.

Light-emitting particles have been a specialty of Bawendi, the Lester Wolf Professor of Chemistry, whose lab has over the years developed new ways of making quantum dots. These nanocrystals, made of semiconductor materials, emit light whose frequency can be precisely tuned by controlling the exact size and composition of the particles.

The key was to develop versions of these quantum dots whose emissions matched the desired short-wave infrared frequencies and were bright enough to then be easily detected through the surrounding skin and muscle tissues. The team succeeded in making particles that are "orders of magnitude better than previous materials, and that allow unprecedented detail in biological imaging," Bruns says. The synthesis of

these new particles was initially described in a paper by graduate student Daniel Franke and others from the Bawendi group in *Nature Communications* last year.

The [quantum dots](#) the team produced are so bright that their emissions can be captured with very short exposure times, he says. This makes it possible to produce not just single images but video that captures details of motion, such as the flow of blood, making it possible to distinguish between veins and arteries.

The new light-emitting [particles](#) are also the first that are bright enough to allow imaging of internal organs in mice that are awake and moving, as opposed to previous methods that required them to be anesthetized, Bruns says. Initial applications would be for preclinical research in animals, as the compounds contain some materials that are unlikely to be approved for use in humans. The researchers are also working on developing versions that would be safer for humans.

The method also relies on the use of a newly developed camera that is highly sensitive to this particular range of short-wave infrared light. The camera is a commercially developed product, Bruns says, but his team was the first customer for the camera's specialized detector, made of indium-gallium-arsenide. Though this camera was developed for research purposes, these frequencies of infrared light are also used as a way of seeing through fog or smoke.

Not only can the new method determine the direction of blood flow, Bruns says, it is detailed enough to track individual blood cells within that flow. "We can track the flow in each and every capillary, at super high speed," he says. "We can get a quantitative measure of flow, and we can do such flow measurements at very high resolution, over large areas."

Such imaging could potentially be used, for example, to study how the blood [flow](#) pattern in a tumor changes as the tumor develops, which might lead to new ways of monitoring disease progression or responsiveness to a drug treatment. "This could give a good indication of how treatments are working that was not possible before," he says.

More information: Next-generation in vivo optical imaging with short-wave infrared quantum dots, *Nature Communications* (2017).

[nature.com/articles/doi:10.1038/s41551-017-0056](https://doi.org/10.1038/s41551-017-0056)

Provided by Massachusetts Institute of Technology

Citation: Quantum dots that emit infrared light open new window for biological imaging (2017, April 10) retrieved 27 April 2024 from <https://phys.org/news/2017-04-quantum-dots-emit-infrared-window.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.