

Scientists publish study on glacial carbon cycle

April 4 2017



Christine Foreman, left, and Heidi Smith, right, in the McMurdo Dry Valleys region of Antarctica. Photo courtesy Christine Foreman. Credit: Montana State University

Two Montana State University researchers have played a major role in

discovering how microbial communities in melting glaciers contribute to the Earth's carbon cycle, a finding that has global implications as the bulk of Earth's glaciers shrink in response to a warming climate.

Heidi Smith, a postdoctoral researcher, and Christine Foreman, associate professor of chemical and biological engineering, both of the Center for Biofilm Engineering in MSU's College of Engineering, were co-authors of a paper published in the prestigious journal *Nature Geoscience*. The paper was published on the journal's website on April 3.

Titled "Microbial formation of labile organic carbon in Antarctic glacial environments," the article was co-authored by researchers at the University of Colorado at Boulder, the U.S. Geological Survey, Stockholm University in Sweden and the Max Planck Institute for Marine Microbiology in Germany.

The paper challenges the prevailing theory that microorganisms found in glacial meltwater primarily consume ancient organic carbon that was once deposited on glacial surfaces and incorporated into ice as [glaciers](#) formed.

"We felt that there was another side to the story," said Smith, the paper's lead author. Smith earned a Ph.D. in ecology and environmental sciences in MSU's Department of Land Resources and Environmental Sciences in 2016, with Foreman as her adviser.

"What we showed for the first time is that a large proportion of the organic carbon is instead coming from photosynthetic bacteria" that are also found in the ice and that become active as the ice melts, Smith said. Like plants, those bacteria absorb carbon dioxide and in turn provide a source of organic matter.

The research team made the discovery after sampling meltwater from a

large stream flowing over the surface of a glacier in the McMurdo Dry Valleys region of Antarctica in 2012.



The Cotton Glacier stream in the McMurdo Dry Valleys region of Antarctica, where MSU researchers Heidi Smith and Christine Foreman sampled glacier runoff in 2012, is shown in this aerial photo. Credit: Christine Foreman

Afterward, Smith spent two months at the Max Planck Institute for Marine Microbiology in Bremen, Germany, with support from the National Science Foundation's flagship interdisciplinary training program, the Integrative Graduate Education and Research Traineeship. There, she worked with colleagues to track how different carbon isotopes moved through the meltwater's ecosystem, allowing the team to determine the carbon's origin and activity.

The researchers ultimately found that the glacial microbes utilized the carbon produced by the photosynthetic bacteria at a greater rate than the older, more complex carbon molecules deposited in the ice, because the bacterial carbon is more "labile," or easily broken down. The labile carbon "is kind of like a Snickers bar," meaning that it's a quick, energizing food source that's most available to the microbes, Smith said.

Moreover, the researchers found that the [photosynthetic bacteria](#) produced roughly four times more carbon than was taken up by the microbes, resulting in an excess of organic carbon being flushed downstream. "The ecological impact of this biologically produced organic carbon on downstream ecosystems will be amplified due to its highly labile nature," Foreman said.

Although individual glacial streams export relatively small amounts of organic carbon, the large mass of glaciers, which cover more than 10 percent of the Earth's surface, means that total glacial runoff is an important source of the material. Marine [organic carbon](#) underpins wide-ranging ecological processes such as the production of phytoplankton,

the foundation of the oceans' food web.

As glaciers increasingly melt and release the organically produced, labile carbon, "we think that marine [microbial communities](#) will be most impacted," Smith said. "We hope this generates more discussion."

In a "News and Views" commentary accompanying the article in *Nature Geoscience*, Elizabeth Kujawinski, a tenured scientist at Woods Hole Oceanographic Institution, called the team's work "an elegant combination" of research methods.

Taken together with another study published in the same issue of *Nature Geoscience*, about microbial carbon cycling in Greenland, Smith's paper "deflates the notion that glacier surfaces are poor hosts for microbial metabolism," according to Kujawinski. The two studies "have established that microbial [carbon](#) cycling on glacier surfaces cannot be ignored," she added.

More information: H. J. Smith et al. Microbial formation of labile organic carbon in Antarctic glacial environments, *Nature Geoscience* (2017). [DOI: 10.1038/ngeo2925](https://doi.org/10.1038/ngeo2925)

Provided by Montana State University

Citation: Scientists publish study on glacial carbon cycle (2017, April 4) retrieved 1 May 2024 from <https://phys.org/news/2017-04-scientists-publish-glacial-carbon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
