

Stabilizing molecule could pave way for lithium-air fuel cell

April 26 2017, by Tom Fleischman

Lithium-oxygen fuel cells boast energy density levels comparable to fossil fuels and are thus seen as a promising candidate for future transportation-related energy needs.

Several roadblocks stand in the way of realizing that vision, however. They include poor rechargeability, reduced efficiency due to high overpotentials (more charge energy than discharge energy) and low specific energy.

Two instabilities contribute to these roadblocks. Much of the previous work done in the lab of Lynden Archer, the James A. Friend Family Distinguished Professor of Engineering in the Robert F. Smith School of Chemical and Biomolecular Engineering (CBE), has centered on one: the nucleation and growth of dendrites from one electrode to the other, which causes short-circuiting, a source of premature cell failure that invariably ends in fires.

It's the other instability – the loss of battery power, also known as capacity fade – that is the focus of the lab's most recent work. Snehashis Choudhury, a doctoral student in the Archer Research Group, has come up with what Archer terms an "ingenious" answer to the problem of capacity fade.

Their work is detailed in "Designer interphases for the lithium-oxygen electrochemical cell," published April 21 in *Science Advances*. Choudhury is co-first author along with Charles Wan '17, a chemical

engineering major.

Capacity fade occurs when the electrolyte, which transports charged ions from the negative electrode (anode) to the positive (cathode), reacts with the electrodes. "It starts to consume the electrodes," Choudhury said. "It forms many insulating products that impede ion transport. Over time, these build up to produce such prohibitive internal cell resistance that finally the battery fades."

The problem: How do you stop one electrolyte-electrode reaction, when it's another necessary reaction between the two – the transfer of ions – that produces power? Choudhury's solution is called an artificial solid-electrolyte interphase (SEI), a material that protects the electrodes while promoting the flow of electrons from one end of the cell to the other.

"Such interphases form naturally in all electrochemical cells ... and their chemo-mechanical stability is critical to the success of the graphite anode in lithium-ion batteries," Archer said. "

Choudhury's approach for creating a functional designer interphase is based on bromide-containing ionic polymers (ionomers) that selectively tether to the lithium anode to form a few-nanometers-thick conductive coating that protects the electrode from degradation and fade. The SEI ionomers display three attributes that allow for increased stability during electrodeposition: protection of the anode against growth of dendrites; reduction-oxidation (redox) mediation, which reduces charge overpotentials; and the formation of a stable interphase with lithium, protecting the metal while promoting ion transport.

One challenge still exists: All research-grade lithium-oxygen electrochemical cells are evaluated using pure oxygen as the active cathode material. For a commercially viable lithium-oxygen (or lithium-air, as it's also known) cell, it would need to pull oxygen out of the air,

and that oxygen also contains other reactive components, such as moisture and carbon dioxide.

If the inefficiencies that limit performance of lithium-oxygen fuel cells can be resolved, the exceptional energy storage options offered by the cell chemistry would be a giant step forward for electrified transportation and a revolutionary advance for autonomous robotics, Archer said.

"It is telling from observations of the most advanced humanoid robots that they are always either tethered to an ultra-long electrical cable or are using something like a loud lawnmower engine to generate energy," Archer said. "Either energy source compares poorly to those living systems have developed—energy storage technologies such as Li-air cells that harness materials from the surroundings promise to close this gap."

Other contributors were Lena Kourkoutis, assistant professor and the Rebecca Q. and James C. Morgan Sesquicentennial Faculty Fellow in applied and engineering physics; CBE doctoral student Wajdi Al Sadat; Sampson Lau, Ph.D. '16; Zhengyuan Tu, doctoral student in materials science and engineering; and Michael Zachman, doctoral student in applied and engineering physics.

Archer noted that Wan and Lau built the electrochemical cell, including designing the cathode configuration, used in their experimentation.

"Charles is an exceptional undergraduate student," Archer said.

"Undergraduates are here principally to focus on getting a first-rate education and historically have had little time to conduct research. But increasingly they are engaging in research, and at a level that in some cases is comparable to our best Ph.D. students."

"I am truly fortunate to have Professor Archer as a mentor," Wan said. "This publication is proof that undergraduates can play a critical role in research if given the chance, something Professor Archer wholeheartedly believes."

More information: Snehashis Choudhury et al. Designer interphases for the lithium-oxygen electrochemical cell, *Science Advances* (2017). DOI: [10.1126/sciadv.1602809](https://doi.org/10.1126/sciadv.1602809)

Provided by Cornell University

Citation: Stabilizing molecule could pave way for lithium-air fuel cell (2017, April 26) retrieved 13 March 2024 from <https://phys.org/news/2017-04-stabilizing-molecule-pave-lithium-air-fuel.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--