

Detecting bloodstains—with an antimalarial compound

May 31 2017



Telling which stain on these paper filters (top row) is from blood (C) rather than coffee (A) or tea (B) is easier with a new compound that reacts with blood and glows in the dark (bottom row, c). Credit: American Chemical Society

As seen on crime shows, investigators use a combination of luminol and



other substances to light up bloodstains at crime scenes. But now, researchers report in ACS' journal *Analytical Chemistry* that combining luminol with artemisinin, a natural peroxide and antimalarial treatment, reduces the risk of false positives compared to the traditional method.

Luminol is often combined with <u>hydrogen peroxide</u> to react with the heme groups in blood, producing a bright blue glow, known as chemiluminescence. This glow allows crime scene technicians to detect blood that has dried on surfaces or to detect blood that someone tried to clean from a surface. While this test has proven beneficial in numerous criminal investigations, it isn't perfect. False positives due to interferences from biomolecules and metal ions, as well as from the breakdown products of hydrogen peroxide, can occur. In an effort to minimize erroneous identifications, Guobao Xu and colleagues investigated the use of a compound called artemisinin, commonly known as a treatment for malaria, with luminol. Artemisinin is more structurally sound and is more resistant to interferences than hydrogen peroxide.

The researchers showed that the luminol-artemisinin combination is more selective than luminol-hydrogen peroxide, resulting in fewer <u>false</u> <u>positives</u> when challenged with components of bleaches and disinfectants, which criminals often use to cover up a misdeed. The researchers found that the method could distinguish blood from coffee, tea and brown sugar stains. They also obtained test results with the new luminol method using a smartphone, which could provide highly accurate on-scene analyses.

More information: Wenyue Gao et al. Artemisinin-Luminol Chemiluminescence for Forensic Bloodstain Detection Using a Smart Phone as a Detector, *Analytical Chemistry* (2017). <u>DOI:</u> <u>10.1021/acs.analchem.7b01000</u>

Abstract



Forensic luminol chemiluminescence test is one of the most sensitive and popular methods for the determination of latent bloodstains. It mainly uses hydrogen peroxide or sodium perborate as coreactants. The easy decomposition of hydrogen peroxide and sodium perborate in the presence of many ions significantly affects the selectivity. Artemisinin is a natural peroxide that is quite stable in the presence of common ions. In the present study, artemisinin has been exploited for the forensic bloodstain chemiluminescence detection for the first time. Using smart phone as cost-effective portable detector, the visual detection of bloodstains has been achieved with a dilution factor of blood up to 100 000. Moreover, this system shows excellent selectivity against many common species. It can well differentiate bloodstains from other stains, such as coffee, brown sugar, and black tea. Both favorable sensitivity and selectivity makes the present method promising in forensic detection.

Provided by American Chemical Society

Citation: Detecting bloodstains—with an antimalarial compound (2017, May 31) retrieved 30 April 2024 from <u>https://phys.org/news/2017-05-bloodstainswith-antimalarial-compound.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.