

# Bumblebee populations higher in Detroit than in some less-urbanized areas; vacant lots could be a factor

May 17 2017

---



Image credit: Wikipedia.

A new study of native bumblebee populations in southeastern Michigan cities found, surprisingly, that Detroit has more of the large-bodied bees than some surrounding, less urbanized locations.

The University of Michigan students who conducted the study suspect that the large amount of vacant or idle land in Detroit may boost the [bumblebee](#) population by providing nesting sites and flowers for food.

Bumblebees belong to the bee genus *Bombus*. In the study, more than 500 individuals from 10 species were identified at 30 sites in southeast Michigan.

"Sites within Detroit had higher *Bombus* abundance and diversity, despite their location in the densest urban landscape," according to the authors—four U-M graduate students and one undergraduate—of a study published online May 17 in the journal *Royal Society Open Science*. "Overall, these results have important implications for conservation of native bee populations and pollination services."

Native bees are critical sources of pollination for agriculture and wild flowering plants. Many [native bees](#) are declining in both abundance and diversity, due to various causes that likely include loss of habitat from human activities. While the effects of large-scale agriculture on native bees are relatively well understood, the effects of urban development are less clear.

To help clarify the role of urbanization, the U-M researchers sampled bumblebees at nature reserves and urban farms and gardens in several southeast Michigan cities—Dexter, Ann Arbor, Ypsilanti, Dearborn and Detroit—with varying degrees of urban development. Each city has a dense urban core surrounded by suburban development.

Bumblebees need to nest in less-disturbed areas with bare ground, tall grass or abandoned tree stumps, making them a good candidate for testing the effects of urban land development. Handheld nets and insect traps were used to capture the bees.

The researchers used geographical information systems data to develop profiles of the land cover types surrounding each of the 30 study sites. As a proxy for [urban development](#), they used the amount of impervious surfaces—buildings, concrete parking structures, asphalt roads, etc.—blanketing the ground.

They found that increased urbanization, as measured by the proportion of impervious surface area, was tied to decreases in both the abundance

and diversity of bumblebees sampled at sites outside Detroit.

But that pattern did not hold in Detroit. There, high urbanization seemed to be correlated with increased bumblebee abundance when compared to less urbanized sites, a finding the authors describe as "an initially unintuitive trend."

In fact, the Detroit site with the highest impervious surface coverage, an urban agriculture demonstration garden in downtown Detroit, had nearly the same number of bumblebees captured as U-M's E.S. George Reserve, a nature preserve near Pinckney with the lowest impervious surface cover proportion in the study.

One possible explanation is that the thousands of vacant residential properties in Detroit help sustain [bumblebee colonies](#), according to the authors. Vacant lots are often less frequently mowed and less likely to be treated with pesticides and herbicides. Therefore, these lots can provide various flowering plants and nesting sites for bees.

The most abundant bumblebee species sampled during the study was the common eastern bumblebee (72.1 percent), followed by the brown-belted bumblebee (11.4 percent) and the two-spotted bumblebee (9.6 percent).

Another unexpected study finding was that urbanization seems to affect female and male bumblebees differently. Those distinct and previously unidentified effects seem to stem from behavioral differences between sexes.

Bumblebee colonies consist of one reproductive female, many female workers that forage for nectar and pollen and bring it back to the colony, and reproductive male drones whose only role is to mate. These different roles lead to distinct movement patterns and nesting needs between the

sexes.

By analyzing males and females separately, the U-M researchers found that observed declines in overall bumblebee abundance and diversity with increasing urbanization were entirely driven by declines in female workers, while male abundance and diversity were unrelated to urbanization.

The student researchers said their results highlight the importance of heterogeneity in urban areas and suggest that urban landscapes can be managed to support native bee conservation.

They thanked the farmers and gardeners who offered their plots for the study. "We are very grateful to everyone who participated. We could not have done this research without their support," said first author Paul Glaum.

**More information:** Paul Glaum et al, Big city: using natural history and land-use history to find significant environmental drivers in bumblebee declines in urban development, *Royal Society Open Science* (2017). [DOI: 10.1098/rsos.170156](https://doi.org/10.1098/rsos.170156)

Provided by University of Michigan

Citation: Bumblebee populations higher in Detroit than in some less-urbanized areas; vacant lots could be a factor (2017, May 17) retrieved 26 April 2024 from <https://phys.org/news/2017-05-bumblebee-populations-higher-detroit-less-urbanized.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.