

Travel distances of juvenile fish key to better conservation

May 16 2017



As part of the largest, most comprehensive study of larval dispersal ever conducted, scientists were able to determine that most of the juvenile clownfish stayed relatively close to home, settling at mean distances of 10-15 kilometers from their natal reefs. Credit: Simon Thorrold, Woods Hole Oceanographic Institution

Marine reserves—sections of the ocean where fishing is prohibited—promote coral reef sustainability by preventing overfishing

and increasing fish abundance and diversity. But to be effective, they need to be sized right, and in a way that accounts for how far juvenile fish travel away from their parents after spawning.

Scientists at the Woods Hole Oceanographic Institution (WHOI), along with researchers from Australia, France, and Saudi Arabia, have successfully measured the dispersal distances of two [coral reef fish](#) species across a 3,000 square mile section of the ocean—an area the size of Yellowstone National Park. The study, published in the May 8, 2017, issue of the journal *Nature Ecology & Evolution*, marks the largest, most comprehensive study of larval dispersal ever conducted and has important implications for the sizing and spacing of [marine reserves](#).

"How far fish will disperse in their lifetimes is critical when you start thinking about how marine reserves should be designed," said Simon Thorrold, co-author of the study and a senior scientist at WHOI. "This is the first time we've been able to measure dispersal distances on spatial scales that are relevant to marine reserves, which means we can now provide data that informs management on optimal spacing and sizing."

Size matters

Marine reserves come in many shapes and sizes. But if a reserve is too small, it can't accommodate enough larvae to sustain populations. And if it's too big, larvae will simply stay within the confines of the reserve without contributing to surrounding fisheries—a critical secondary role marine reserves need to play to improve fisheries management.

To get a read on fish dispersal in the past, scientists relied on population genetics approaches that lacked the power to measure dispersal over space and time scales relevant to protected areas of the ocean. More recently, ecologists have turned to computer-generated models of water currents to track particles through virtual oceans. According to Thorrold,

this approach also has limitations since there was no way to verify the accuracy of the models. "The software can generate a lot of cool-looking graphs, but it was impossible to test the skill of those models in any real way."

An empirical approach

To overcome these limitations, Thorrold and his colleagues took direct measurements of dispersal distances in the field. They collected DNA samples from thousands of adult and juvenile clownfish and butterflyfish throughout Kimbe Bay, Papua New Guinea, in 2009 and 2011. The entire sampling process occurred underwater, with the 30-person science team spending thousands of man-hours on SCUBA over several weeks in the field each year.



How far fish will disperse in their lifetimes is critical when you start thinking about how marine reserves should be designed, according to WHOI biologist

Simon Thorrold, co-author of the new study. The research team found that butterflyfish dispersed further than clownfish, averaging distances of 43-64 kilometers before settling into their new habitats. Credit: Simon Thorrold, Woods Hole Oceanographic Institution)

When the scientists returned to the lab, they used DNA parentage analysis, a sequencing technique that allowed them to match the juveniles up with their parents based on the DNA samples and spawning and settlement location data. From that, they were able to determine that most of the juvenile clownfish stayed relatively close to home, settling at mean distances of 10-15 kilometers from their natal reefs. The butterflyfish dispersed further, averaging distances of 43-64 kilometers before settling into their new habitats.

"Since we knew the respective locations of the adults and babies, we were able to come up with the exact linear distances that the larvae had dispersed. We're no longer talking about estimates," said Thorrold.

Benefits beyond design

In addition to helping inform the design of protected areas, the measurements can help to test the ability of reserves to perform key conservation functions. For example, one way a marine reserve network may improve fish population sustainability is through the so-called "rescue effect." In theory, if fish in a reserve suffer catastrophic mortality, the reserve can be repopulated by larvae from other reserves within the network. Thorrold and colleagues were able to track larvae from one reserve to another in the study area, confirming that rescue effect is likely to occur in real-world reserve networks.

The dispersal measurements could also allow fisheries managers to

monitor the effectiveness of existing reserves, helping answer the question of whether or not a particular reserve is contributing to fish populations beyond its boundaries. This, according to Thorrold, has been a big unknown.

"If you can trace larvae from one reserve to a place that's fished, you can come up with a direct measure of how many fish the reserve is contributing to exploited populations beyond the reserve," he said. "This helps when trying to convince fishermen that networks of marine reserves are a good management tool."

Future work

According to Thorrold, as coral reef seascapes continue to face pressure from man-made stressors, marine reserves will continue to serve as an important conservation management tool. As such, it will become increasingly important to be able to provide direct measurements of larval dispersal, and find ways to apply the information to other regions of the ocean.

"The next thing we are working on is developing a coupled bio-physical model of the area that will allow us to take the results from this study and generalize them to other coral reef seascapes around the world," he said. "Limited resources for ocean management, particularly in the developing world, means that we need to maximize the chances of successful conservation outcomes from these efforts. These types of scientific insights will be critical for ongoing efforts to promote resilience of coral reef ecosystems in the face of human exploitation and climate change."

Provided by Woods Hole Oceanographic Institution

Citation: Travel distances of juvenile fish key to better conservation (2017, May 16) retrieved 13 March 2024 from <https://phys.org/news/2017-05-distances-juvenile-fish-key.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.