

Imaging mRNA right where it is made—at the site of translation

May 3 2017





Researchers use fluorescence probes to monitor increases in mRNA-ribosome interaction levels for a gene associated with iron storage in response to iron (right panels). Scale bar = $20 \mu m$. Credit: American Chemical Society

Think of life as a house: if DNA molecules are blueprints, then messenger RNAs (mRNAs) are orders, describing the required parts (proteins) and when they should arrive. But putting in many orders doesn't always mean you'll get all of the parts on time—maybe there's a delay with your vendor or delivery service. Similarly, mRNA levels alone do not dictate protein levels. Today in *ACS Central Science*, researchers report a method to address that issue.

David Tirrell, Kelly Burke and Katie Antilla note that in order to better understand how genes are regulated, one needs to see the mRNA when it is at the site of protein synthesis. Using fluorescence probes, the researchers designed a technique that shows mRNA when it comes in contact with giant protein synthesizing machines called ribosomes. They used this method to record the synthesis of proteins and to measure cellular responses to iron. Unlike previous methods, their tool works without the need to engineer an mRNA of interest.

Tirrell notes that the method is applicable to essentially any type of RNA, and could be modified to visualize other types of interactions in the cell.

More information: *ACS Central Science* (2017). pubs.acs.org/doi/full/10.1021/acscentsci.7b00048



Provided by American Chemical Society

Citation: Imaging mRNA right where it is made—at the site of translation (2017, May 3) retrieved 1 May 2024 from <u>https://phys.org/news/2017-05-imaging-mrna-madeat-site.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.