

## **Micro delivery service for fertilizers**

May 22 2017



Credit: Wiley

Plants can absorb nutrients through their leaves as well as their roots. However, foliar fertilization over an extended period is difficult. In the journal *Angewandte Chemie*, German researchers have now introduced an efficient delivery system for micronutrients based on biohybrid microgels. Special peptides anchor the "microcontainers" onto the leaf surface while binding sites inside ensure gradual release of the "cargo".

Foliar fertilization is already commonly used in areas such as viniculture,



when the leaves on the vines turn yellow due to a mineral deficiency. Yet, despite the use of detergents, adhesives, and humectants, controlled delivery of nutrients through foliar fertilization over several weeks is nearly impossible to achieve. Up to 80% of the nutrients are washed away, winding up in the soil and being converted into forms that the plant cannot use. In addition, they can be washed into bodies of water and cause environmental problems. An additional problem is that strong sunlight evaporates the water out of the applied fertilizer solution. This results in a high salt concentration that draws water out of the leaf and causes burn damage.

A team from DWI-Leibniz Institute for Interactive Materials in Aachen, RWTH Aachen University, and the University of Bonn has now developed a foliar fertilization system based on biocompatible microgels that adhere selectively to leaves for a long period and slowly deliver nutrients in a controlled fashion. Microgels are tiny particles of crosslinked macromolecules that can bind water and other molecules, such as fertilizers very efficiently.

Led by Ulrich Schwaneberg and Andrij Pich, the researchers equipped the interiors of <u>gel particles</u> with binding sites modeled on the ironbinding proteins of bacteria. These ensure that the iron ions are released slowly. The microgels are loaded with an iron-containing solution at a pH of 3. When the pH rises to 7, the microgels shrink, releasing water and binding the iron.

The surface of the gel particles is equipped with anchor peptides from <u>lactic acid bacteria</u>. These bind securely to leaf surfaces to hinder rinsing away of the microgels. The <u>water</u> in the gel provides an aqueous microenvironment that allows the iron to diffuse into the leaves. Yellow leaves of iron-deficient cucumber plants rapidly turned green in spots where the new foliar fertilizer was applied.



By incorporating different binding sites, the microgel "containers" can be loaded with a multitude of other metal ions or agents. A controlled delivery of agents as required would minimize the applied quantities as well as the release of fertilizers and pesticides into the environment. Low production costs, high levels of loading, easy application, and adjustable adhesive properties should make broad industrial applications possible. The goal is to make self-regulating delivery systems for sustainable agriculture.

**More information:** Richard A. Meurer et al, Biofunctional Microgel-Based Fertilizers for Controlled Foliar Delivery of Nutrients to Plants, *Angewandte Chemie International Edition* (2017). <u>DOI:</u> <u>10.1002/anie.201701620</u>

Provided by Wiley

Citation: Micro delivery service for fertilizers (2017, May 22) retrieved 5 May 2024 from <u>https://phys.org/news/2017-05-micro-delivery-fertilizers.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.