

Neonicotinoid pesticide reduces egg development in wild bumblebee queens

May 3 2017



Bombus Terrestris Bee. Credit: Royal Holloway, University of London

New research published in the journal Proceedings of the Royal Society B



has found that wild bumblebee queens are less able to develop their ovaries when exposed to a common neonicotinoid pesticide

The research was conducted by Dr Gemma Baron , Professor Mark Brown of Royal Holloway, University of London and Professor Nigel Raine, (now based at the University of Guelph).

The study investigated the <u>impact</u> of exposure to field-realistic levels of a neonicotinoid insecticide (thiamethoxam) on the feeding behaviour and ovary development of four <u>species</u> of bumblebee <u>queen</u>.

Reduced ovary development across bee species

Lead author, Dr Baron said, "We consistently found that neonicotinoid exposure, at levels mimicking exposure that queens could experience in agricultural landscapes, resulted in reduced ovary development in queens of all four species we tested."

She continued, "Impacts of <u>neonicotinoid</u> exposure on feeding behaviour were species-specific, with two out of four species eating less artificial nectar when exposed to the pesticide. These impacts are likely to reduce the success of bumblebee queens in the spring, with knock-on effects for bee populations later in the year" added Baron.

As the first to examine the impacts of these chemicals across multiple bumblebee species, this study is an important contribution to understanding the potential costs of using this class of insecticides.

Dr Baron explained, "Previous studies have focused on a single bumblebee species and examined impacts in workers and established colonies. Bumblebee populations rely on spring queens to succeed, and by looking at the impacts of thiamethoxam on multiple species of spring queens, we have gained a step-change in our understanding".



Impact on EU pesticide regulations

Thiamethoxam is currently under an EU moratorium, along with two other neonicotinoids.

"The EU, and other countries, is weighing the benefits of using neonicotinoids to protect crops, against real concerns about how such pesticides can cause unintended harm to pollinators and other beneficial insects." added Professor Raine. "This work on wild <u>bumblebee</u> queens adds significantly to the evidence base on which to make such important policy decisions".

Professor Brown commented that. "Our work provides a major step forward in understanding the general and species-specific impacts of neonicotinoids on wild bees. Future studies across different species are likely to demonstrate further variation in the impact of neonicotinoids, and conducting such studies needs to be a priority for scientists and governments."

More information: Gemma L. Baron et al. General and speciesspecific impacts of a neonicotinoid insecticide on the ovary development and feeding of wild bumblebee queens, *Proceedings of the Royal Society B: Biological Sciences* (2017). DOI: 10.1098/rspb.2017.0123

Provided by Royal Holloway, University of London

Citation: Neonicotinoid pesticide reduces egg development in wild bumblebee queens (2017, May 3) retrieved 2 May 2024 from https://phys.org/news/2017-05-neonicotinoid-pesticide-egg-wild-bumblebee.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.