


# Preparations for a new season of physics at the Large Hadron Collider

May 16 2017



One of the first proton-proton collisions seen by the ALICE experiment in 2017, on May 13, during the LHC beam commissioning phase. ALICE used these first collisions to fine-tune its equipment and get ready for the new physics season of LHC. Credit: CERN


Last week, the detectors of the Large Hadron Collider (LHC) witnessed their first collisions of 2017. These test collisions were not for physics research, instead they were produced as part of the process of restarting

the LHC. But have patience, data taking for physics will start in another few days.

Since [particles began circulating in the large ring once more](#), the LHC's operators have been testing and adjusting 24 hours a day to turn the LHC into a veritable [collision](#) factory. Their work involves forming trains of bunches, building them up over the next few weeks to several hundred and then several thousand bunches per beam.

To establish this production line of particles, all of the accelerator's systems must be perfectly adjusted. The LHC is an extremely complex machine comprising thousands of subsystems and it takes weeks to adjust them all.


The first particles circulated on 29 April 2017 and, soon after, the operators started work on their long list of adjustments. They tested the radiofrequency system, which accelerates the particles. They brought the [beam energy](#) up to its operating value of 6.5 TeV. They tested the beam dump system, which ejects the particles into a block of graphite if required. They tested and aligned all the collimators – jaw-like devices that close around the beam to absorb stray [particles](#). They carried out proton bunch ramp and squeeze cycles. Finally, they performed fine adjustments of the hundreds of corrector magnets, adjusting the trajectory of the [beam](#) to a precision of one micron at the collision points.



This image shows a beam splash, as observed by the ATLAS experiment on 29 April, the day of the LHC restart. The beam splashes are generated by aiming beams at the collimators near to the experiments, in this case 140 metres from the ATLAS interaction point. Once the LHC is back in operation, the experiments use the beam splashes to synchronise their sub-detectors with the accelerator's clock. Credit: CERN

Last Wednesday, they started to collide the beams to be able to adjust the interaction points at the heart of the experiments. This step is carried out with so-called "pilot" beams, containing fewer than ten bunches and fewer protons than during the physics runs. These first collisions also allow the experiments to adjust their detectors.

In the coming days, the operators will continue to adjust and align the equipment. Once all of these steps are complete, they will be able to announce "stable beams", the long-awaited signal for the start of the new data-taking season for the [experiments](#).



A beam splash, as observed by the CMS experiment on 29 April. In contrast to proton-proton collisions where the particles come from the center of the detector, in splash events particles traverse the detector horizontally, from one side to the other. Credit: CERN

Provided by CERN

Citation: Preparations for a new season of physics at the Large Hadron Collider (2017, May 16)  
retrieved 23 April 2024 from

<https://phys.org/news/2017-05-season-physics-large-hadron-collider.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.