


Rising temperatures threaten stability of Tibetan alpine grasslands

May 10 2017



Researchers studied the effects of rising temperatures and varying rainfall on Tibetan Plateau grasslands at this experimental site at the Haibei Alpine Grassland Ecosystem Research Station of the Chinese Academy of Sciences. Credit: Xian Yang and Qianna Xu

A warming climate could affect the stability of alpine grasslands in

Asia's Tibetan Plateau, threatening the ability of farmers and herders to maintain the animals that are key to their existence, and potentially upsetting the ecology of an area in which important regional river systems originate, says a new study by researchers in China and the United States.

Though temperature changes could destabilize the fragile ecosystem of the area, variations in rainfall appear to have no similar effect. The study involved varying two factors likely to change with a warming climate - temperature and rainfall - in test plots over a five-year period. The project is believed to be the first to simultaneously examine the effects of temperature and rainfall changes on [ecosystem stability](#).

"We were concerned about the variability of the total community plant cover over time," said Lin Jiang, a professor in the Georgia Tech School of Biological Sciences. "Significant warming could reduce the stability of the grasslands, which would increase the variability of plant biomass production that could be a significant issue for people living in the region. We believe the effects of climate change could be particularly dramatic in this area."

The research, conducted by scientists from Peking University, the Chinese Academy of Sciences and the Georgia Institute of Technology, was scheduled to be published May 10 in the journal *Nature Communications*. The research was supported by the National Basic Research Program of China, the National Nature Science Foundation of China and the U.S. National Science Foundation.

The Tibetan Plateau is an area of about 2.5 million square kilometers in which summertime high temperatures seldom rise above 25 degrees Celsius and nighttime temperatures could drop below freezing even in the summer. Because of the altitude, temperature extremes and high winds, more than two-thirds of the Plateau is grassland used for grazing

yak, sheep and other animals. About 9.8 million people live in the area, which is also the source for several of Asia's major river systems.



The Tibetan Plateau is an area of about 2.5 million square kilometers in which summertime high temperatures seldom rise above 25 degrees Celsius and nighttime temperatures could drop below freezing even in the summer. Credit: Xian Yang and Qianna Xu

"Our results suggest that under a warmer climate, the ecosystem would provide less forage production in drought years, and more biomass production in wet years - which is undesirable," said Jin-Sheng He, a

professor in the Department of Ecology, College of Urban and Environmental Sciences at Peking University. "Reduced plant production temporal stability could mean that this alpine ecosystem may not be able to provide stable forage for the livestock that local people rely on. Reduced stability may also have consequences for other ecosystem services, such as climate regulation and water conservation."

The researchers found that the stability of the grasslands was affected not by the richness of plant species, but by the effects on dominant species and the asynchrony of the species.

"We found that climate warming lowers stability through increasing species synchrony in which the biomass of a few dominant species increased while that of most rare species declined," said He. "That indicates the alpine grasslands that have well adapted to cold environments owing to their long-term evolutionary history may be jeopardized in the future."

Experimentally, the researchers created test plots in which some were heated to two degrees Celsius above the surrounding grasslands. At the same time, the researchers varied the amount of rainfall onto the plots, with some sections receiving 50 percent more water, and others receiving 50 percent less. There were also control sections in which temperature and rainfall were not adjusted. Each of the six conditions were replicated six times, for a total of 36 test plots.

Over a period of five years, the researchers studied the growth of different grass species by weighing the biomass production from the different test plots. The research was done at the Haibei Alpine Grassland Ecosystem Research Station of the Chinese Academy of Sciences.



Because of the altitude, temperature extremes and high winds, more than two-thirds of the Tibetan Plateau is grassland used for grazing yak, sheep and other animals. Rising temperatures could affect the area's ability to sustain these animals. Credit: Xian Yang and Qianna Xu

Jiang was surprised that the dramatic variations in rainfall didn't affect the grass species, and hadn't expected much impact from the temperature change, which translates to about 3.6 degree Fahrenheit. "The plants appear to be able to tolerate significant variations in the amount of water available," he said.

While the Tibetan Plateau is unique for its size and high average altitude, there are other areas of the world with similar conditions. "If these

findings can be generalized to other alpine ecosystems, we may need to be concerned about large variations in [biomass production](#) in these other areas, as well," Jiang added.

The study adds to knowledge about the Tibetan plateau, which has not been well studied because of its geographical isolation and harsh climate.

"The Tibetan plateau is sometimes called the third pole because there are so many high mountains and so much of the area is covered with snow and ice," said Jiang. "Ecologically, it is a very important region, but relatively few ecological studies have ever been done there."

More information: Zhiyuan Ma, Huiying Liu, Zhaorong Mi, Zhenhua Zhang, Yongui Wang, Wei Xu, Lin Jiang and Jin-Sheng He, "Climate warming reduces the temporal stability of community biomass production," *Nature Communications*, 2017. [DOI: 10.1038/NCOMMS15378](#)

Provided by Georgia Institute of Technology

Citation: Rising temperatures threaten stability of Tibetan alpine grasslands (2017, May 10) retrieved 27 April 2024 from <https://phys.org/news/2017-05-temperatures-threaten-stability-tibetan-alpine.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
