

# 'Tiny clocks' crystallize understanding of meteorite crashes

May 26 2017

---



This rocky outcrop at Sudbury is where the crystals of baddeleyite came from -- crystals that are now being used in a new technology to help date when meteorite strikes took place. Credit: Desmond Moser/Western University

Almost two billion years ago, a 10-kilometre-wide chunk of space slammed down into rock near what is now the city of Sudbury. Now, scientists from Western University and the University of Portsmouth are marrying details of that meteorite impact with technology that measures surrounding crystal fragments as a way to date other ancient meteorite strikes.

The pioneering technique is helping add context and insight into the age of meteor impacts. And ultimately, it provides new clues into the beginnings of life on this planet and others, said Desmond (Des) Moser, associate professor in the Departments of Earth Sciences and Geography at Western.

"The underlying theme is, when did life begin? We know that it couldn't happen as long as the surface was being periodically vaporized by [meteorite](#) strikes during the solar system's early years and youth—so if we can figure out when those strikes stopped, we can then understand a bit more about how we got here, and when."

In this instance, researchers have been able to use new imaging techniques to measure the atomic nanostructure of ancient crystals at impact locations, using the 150-kilometre-wide crater at Sudbury as a test site.

Shock waves from that [meteorite impact](#) deformed the minerals that made up the rock beneath the crater, including small, tough crystals that contain trace amounts of radioactive uranium and lead. "These can be used as tiny clocks that are the basis for our [geologic time scale](#)," Moser said. "But because these crystals are a banged-up mess, conventional methods won't help in extracting age data from them."

An international team using specialized instruments at Western's Zircon and Accessory Phase Laboratory (ZAPLab) and a new instrument called

the atom probe, at CAMECA Laboratories in the US, have made that job easier. With the probe, researchers are able to slice and lift out tiny pieces of crystal baddeleyite which is common in terrestrial, Martian and lunar rocks and meteorites.

Then Moser's team—including researcher Lee White and co-supervisor James Darling of the University of Portsmouth—measured the deformation in the crystals after sharpening and polishing the pieces into extremely fine needles, then evaporated and identified the atoms and their isotopes layer by layer. The result is a 3D model of the atoms and their positions.

"Using the [atom probe](#) to go from the rock to the crystal to its atomic level is like zooming in with the ultimate Google Earth," Moser says. This atomic-scale approach holds great potential in establishing a more accurate chronology of the formation and evolution of planetary crusts.

The team's findings are published in the journal *Nature Communications*.

**More information:** L. F. White et al, Atomic-scale age resolution of planetary events, *Nature Communications* (2017). [DOI: 10.1038/ncomms15597](#)

Provided by University of Western Ontario

Citation: 'Tiny clocks' crystallize understanding of meteorite crashes (2017, May 26) retrieved 3 May 2024 from <https://phys.org/news/2017-05-tiny-clocks-crystallize-meteorite.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---