

# Arp 299: Galactic Goulash

June 26 2017

---



Credit: X-ray: NASA/CXC/Univ of Crete/K. Anastasopoulou et al,

NASA/NuSTAR/GSFC/A. Ptak et al; Optical: NASA/STScI

What would happen if you took two galaxies and mixed them together over millions of years? A new image including data from NASA's Chandra X-ray Observatory reveals the cosmic culinary outcome.

Arp 299 is a system located about 140 million light years from Earth. It contains two galaxies that are merging, creating a partially blended mix of stars from each galaxy in the process.

However, this stellar mix is not the only ingredient. New data from Chandra reveals 25 bright X-ray sources sprinkled throughout the Arp 299 concoction. Fourteen of these sources are such strong emitters of X-rays that astronomers categorize them as "ultra-luminous X-ray sources," or ULXs.

These ULXs are found embedded in regions where stars are currently forming at a rapid rate. Most likely, the ULXs are binary systems where a neutron star or black hole is pulling matter away from a companion star that is much more massive than the Sun. These [double star systems](#) are called high-mass X-ray binaries.

Such a loaded buffet of high-mass X-ray binaries is rare, but Arp 299 is one of the most powerful star-forming galaxies in the nearby Universe. This is due at least in part to the merger of the two galaxies, which has triggered waves of star formation. The formation of high-mass X-ray binaries is a natural consequence of such blossoming star birth as some of the young massive [stars](#), which often form in pairs, evolve into these systems.

This new composite image of Arp 299 contains X-ray data from

Chandra (pink), higher-energy X-ray data from NuSTAR (purple), and optical data from the Hubble Space Telescope (white and faint brown). Arp 299 also emits copious amounts of infrared light that has been detected by observatories such as NASA's Spitzer Space Telescope, but those data are not included in this composite.

The infrared and X-ray emission of the galaxy is remarkably similar to that of galaxies found in the very distant Universe, offering an opportunity to study a relatively nearby analog of these distant objects. A higher rate of galaxy collisions occurred when the universe was young, but these objects are difficult to study directly because they are located at colossal distances.

The Chandra data also reveal diffuse X-ray emission from hot gas distributed throughout Arp 299. Scientists think the high rate of supernovas, another common trait of star-forming [galaxies](#), has expelled much of this hot gas out of the center of the system.

A paper describing these results appeared in the August 21st, 2016 issue of the *Monthly Notices of the Royal Astronomical Society*.

**More information:** K. Anastasopoulou et al. A deepChandraobservation of the interacting star-forming galaxy Arp 299, *Monthly Notices of the Royal Astronomical Society* (2016). [DOI: 10.1093/mnras/stw1200](#) , *Arxiv*: [arxiv.org/abs/1605.07001](#)

Provided by Chandra X-ray Center

Citation: Arp 299: Galactic Goulash (2017, June 26) retrieved 19 April 2024 from <https://phys.org/news/2017-06-arp-galactic-goulash.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.