

## **Birds' feathers reveal their winter diet**

June 21 2017



Bobolinks molt while in South America for the winter, and the new feathers they grow there reflect their diet. Credit: R.M. Jensen

Influences outside the breeding season can matter a lot for the population health of migratory birds, but it's tough to track what happens once species scatter across South America for the winter months. A study from *The Condor: Ornithological Applications* tries a new approach



for determining what declining migratory grassland birds called Bobolinks eat after they head south for the winter—analyzing the carbon compounds in their plumage, which are determined by the types of plants the birds consume while growing their feathers during their winter molt.

Thanks to a quirk of photosynthesis, rice contains a different ratio of carbon isotopes than most of the native grasses in South America where Bobolinks winter. Rosalind Renfrew of the Vermont Center for Ecostudies and her colleagues took advantage of this, collecting feather samples from wintering Bobolinks in a rice-producing region and a grassland region and from breeding Bobolinks in North America. When they analyzed the feathers' isotopes ratios, the results from South America confirmed that isotopes in Bobolinks' feathers reflected the differences in their diets between regions with and without rice production. The samples taken in North America showed that the winter diet of most individuals was weighted more toward non-rice material, but that rice consumption was highest late in the <u>winter</u>, when rice is nearing harvest and the <u>birds</u> are preparing for their northbound migration.

Rice could be beneficial by providing the birds with needed calories as they prepare for their journey north, but it could also increase Bobolinks' exposure to pesticides and threats from farmers who see them as pests. According to Renfrew and her colleagues, maintaining native grasslands, encouraging integrated pest management programs to reduce toxic pesticide applications, and compensating farmers for crops lost to feeding birds all would be helpful.

"The time spent coordinating the field work for this study may well have been greater than the time spent collecting the data," says Renfrew. "It was truly a team effort, and the assistance we received from our partners was absolutely essential, especially in South America. Aves Argentinas



and the Museo de Historia Natural de Noel Kempff Mercado provided priceless logistical support, and this study could not have happened without them. Some of the same partners have provided input on a Bobolink Conservation Plan that lays out actions to address threats to grassland birds in North and South America, based on results from this and other studies."

"As Bobolink populations continue to decline, Renfrew and her colleagues use state-of-the-art isotope analysis techniques to assess the Bobolink's diet on its South American wintering grounds," according to John McCracken of Bird Studies Canada, an expert on grassland bird conservation who was not involved with the study. "The authors conclude that rice may have negative effects on Bobolinks, owing to its relatively low nutritional quality and from exposure to insecticides."

**More information:** "Winter diet of Bobolink, a long-distance migratory grassland bird, inferred from feather isotopes" *The Condor: Ornithological Applications*, 2017.

Provided by American Ornithological Society

Citation: Birds' feathers reveal their winter diet (2017, June 21) retrieved 2 May 2024 from <u>https://phys.org/news/2017-06-birds-feathers-reveal-winter-diet.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.