

Charred flowers and the fossil record

June 13 2017




Figure 1 from Hudspith and Belcher: Grayscale photographs of post-burn

residues containing char and ash from flowers and associated vegetative material. A: Spikes of *Anemopsis californica*. B: Spadix of *Spathiphyllum wallisii* flowers. C: *Laurus nobilis* flowers clustering in leaf axils (see arrows). D: Differential charcoal production of small isolated charred *Cinnamomum camphora* flowers compared to ashed leaves, tested under the same conditions. E: Intact charred Schisandraceae: *Calycanthus occidentalis* flowers; even detached tepals remain as char post-burn (see arrows). F: Intact charred Schisandraceae: *Illicium cf. henryi* flower. Scale bar in all images is 20 mm. Credit: Geology, Victoria A. Hudspith and Claire M. Belcher

One of the main types of fossil used to understand the first flowering plants (angiosperms) are charred flowers. These charcoals were produced in ancient wildfires, and they provide some evidence for the types of plants that grew millions of years ago. However, when fires burn they not only produce charcoal, but they also destroy it.

This has led scientists to consider whether some types of [flowers](#) in Earth's past were more likely to be destroyed during fires, rather than preserved as fossil charcoals in rocks. In their study published this week in *Geology*, Victoria Hudspith and Claire Belcher show that different types of plants caused fires to burn differently (some hotter and some cooler) and that the different shapes and forms of certain flowers made them more likely to be entirely burned away by [fire](#) and turned to ash, whereas other types were more likely to remain as [charcoal](#).

This is important because if scientists are trying to use charred flowers to understand the diversity of the earliest angiosperms then they also need to consider that some flowers may have been turned to ash in these ancient fires and not preserved as charcoal at all. In other words, paleontologists must now consider that the charcoal fossil record of flowers is unlikely to preserve all types of flower equally, and as a result, they may be missing information about the early evolutionary history of

angiosperms.

Provided by Geological Society of America

Citation: Charred flowers and the fossil record (2017, June 13) retrieved 1 May 2024 from
<https://phys.org/news/2017-06-charred-fossil.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.