

# New 3-D model predicts best planting practices for farmers

June 23 2017

---



Sugarcane planted in with traditional spacing (pictured here) is better for yields but may be worse for plants and soil quality. Credit: Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign

As farmers survey their fields this summer, several questions come to


mind: How many plants germinated per acre? How does altering row spacing affect my yields? Does it make a difference if I plant my rows north to south or east to west? Now a computer model can answer these questions by comparing billions of virtual fields with different planting densities, row spacings, and orientations.

The University of Illinois and the Partner Institute for Computational Biology in Shanghai developed this computer model to predict the yield of different crop cultivars in a multitude of planting conditions. Published in *BioEnergy-Research*, the model depicts the growth of 3D [plants](#), incorporating models of the biochemical and biophysical processes that underlie productivity.

Teaming up with the University of Sao Paulo in Brazil, they used the model to address a question for sugarcane producers: How much yield might be sacrificed to take advantage of a possible conservation planting technique?

"Current sugarcane harvesters cut a single row at a time, which is time-consuming and leads to damage of the crop stands," said author Steve Long, Gutsell Endowed Professor of Plant Biology and Crop Sciences at the Carl R. Woese Institute for Genomic Biology. "This could be solved if the crop was planted in double rows with gaps between the double rows. But plants in double rows will shade each other more, causing a potential loss of profitability."

# Double row spacing better for sugarcane plants, soil but sacrifices up to 10% yield



Double row spacing is better for sugarcane plants, soil but sacrifices up to 10% of yield. Credit: Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign

The model found that double-row spacing costs about 10% of productivity compared to traditional row spacing; however, this loss can

be reduced to just 2% by choosing cultivars with more horizontal leaves planted in a north-south orientation.

"This model could be applied to other [crops](#) to predict optimal planting designs for specific environments," said Yu Wang, a postdoctoral researcher at Illinois who led the study. "It could also be used in reverse to predict the potential outcome for a field."

The authors predict this [model](#) will be especially useful when robotic planting becomes more commonplace, which will allow for many more planting permutations.

The paper "Development of a Three-Dimensional Ray-Tracing Model of Sugarcane Canopy Photosynthesis and Its Application in Assessing Impacts of Varied Row Spacing" is published by *BioEnergy Research*.

**More information:** Yu Wang et al, Development of a Three-Dimensional Ray-Tracing Model of Sugarcane Canopy Photosynthesis and Its Application in Assessing Impacts of Varied Row Spacing, *BioEnergy Research* (2017). [DOI: 10.1007/s12155-017-9823-x](https://doi.org/10.1007/s12155-017-9823-x)

Provided by University of Illinois at Urbana-Champaign

Citation: New 3-D model predicts best planting practices for farmers (2017, June 23) retrieved 25 April 2024 from <https://phys.org/news/2017-06-d-farmers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.