


The first low-cost system for splitting carbon dioxide

June 5 2017



Ball-and-stick model of carbon dioxide. Credit: Wikipedia

Using Earth-abundant materials, EPFL scientists have built the first low-cost system for splitting CO₂ into CO, a reaction necessary for turning renewable energy into fuel.

The future of clean energy depends on our ability to efficiently store

energy from renewable sources and use it later. A popular way to do this is to electrolyze carbon dioxide to carbon monoxide, which is then mixed with hydrogen to produce liquid hydrocarbons like gasoline or kerosene that can be used as fuel. However, we currently lack efficient and Earth-abundant catalysts for the initial splitting of CO₂ into CO and oxygen, which makes the move into [renewable energy](#) expensive and prohibitive. EPFL scientists have now developed an Earth-abundant catalyst based on copper-oxide nanowires modified with [tin oxide](#). The system can split CO₂ with an efficiency of 13.4%. The work is published in *Nature Energy*, and can help worldwide efforts to synthetically produce carbon-based fuels from CO₂ and water.

The research was carried out by the lab of Michael Grätzel at EPFL. Grätzel is known worldwide for the invention of the first ever [dye-sensitized solar cells](#) (or "Grätzel cells"). The catalyst, developed by PhD student Marcel Schreier and postdoc Jingshan Luo, is made by depositing an atomic layer of tin oxide on copper oxide nanowires. By using such Earth-abundant materials, the design keeps the cost of the catalyst low while significantly increasing the yield of CO, as opposed to the other products that are generated from CO₂ electrocatalysis.

The catalyst was integrated into a CO₂ electrolysis system and linked to a triple-junction solar cell (GaInP/GaInAs/Ge) to make a CO₂ electrolyzer. The system uses the catalyst as a bifunctional electrode that both reduces CO₂ into CO and produces oxygen through what is known as the "oxygen evolution" reaction. The two products are separated with a bipolar membrane.

Using solar [energy](#), the system was able to selectively convert CO₂ to CO with an efficiency of 13.4%, and do so with a Faradaic efficiency up to 90%—this describes how efficiently electrical charge is transferred in an electrocatalysis system like the one developed here. "The work sets a new benchmark for solar-driven CO₂ reduction," says Luo.

"This is the first time that such a bi-functional and low-cost [catalyst](#) is demonstrated," adds Schreier. "Very few catalysts—except expensive ones, like gold and silver—can selectively transform CO₂ to CO in water, which is crucial for industrial applications."

More information: Marcel Schreier, Florent Héroguel, Ludmilla Steier, Shahzada Ahmad, Jeremy S. Luterbacher, Matthew T. Mayer, Jingshan Luo, Michael Grätzel. Solar conversion of CO₂ to CO using Earth-abundant electrocatalysts prepared by atomic layer modification of CuO. *Nature Energy* 2, 17087 (05 June 2017).
[nature.com/articles/doi:10.1038/nenergy.2017.87](https://www.nature.com/articles/doi:10.1038/nenergy.2017.87)

Provided by Ecole Polytechnique Federale de Lausanne

Citation: The first low-cost system for splitting carbon dioxide (2017, June 5) retrieved 25 April 2024 from <https://phys.org/news/2017-06-low-cost-carbon-dioxide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.