


# Meltwater lakes under the Antarctic Ice Sheet accelerated glacial retreat in the Earth's past

June 1 2017

---



A satellite image of Antarctica. Credit: USGS, via Wikipedia, Public Domain

During the last glacial period - when the ice in the Antarctic was far thicker and extended further offshore than it does today - it has been speculated that subglacial lakes existed beneath it. An international team of researchers has now successfully sampled the metre-thick sediment layers left behind by these lakes contemporary on the seafloor. This is the outcome of a study by Gerhard Kuhn and colleagues, which was published today (1 June 2017) in the journal *Nature Communications*.

Hundreds of subglacial lakes currently exist beneath the Antarctic Ice Sheet, with Lake Vostok being the largest and best known. The challenges involved in exploring these lake systems, which have remained enclosed for thousands of years, are enormous. For instance, scientists must adhere to strict environmental restrictions so as to avoid polluting the lakes, which involves considerable resources. To date, Russian researchers have only collected samples by drilling into the surface ice of Lake Vostok and testing the water that poured out of the borehole before promptly refreezing.

Now a team of researchers led by marine geologist Dr Gerhard Kuhn from the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI) have provided unequivocal evidence, in the form of pore water analyses, for the presence of a former subglacial lake on the Antarctic continental shelf. During expeditions to the Amundsen Sea with the Research Icebreaker Polarstern in 2006 and 2010, AWI researchers and their international colleagues collected [sediment](#) cores that they now confirm are from subglacial lakes. "The cores, which are up to ten metres long, were collected at a water depth of 750 metres. The [lake sediments](#) are currently buried under a four-metre thick layer of marine sediment on the seafloor," Kuhn reports. They were retrieved from valleys on the ocean floor that were situated under the Antarctic Ice Sheet in the Earth's distant past. "We have now verified that, during the last [glacial period](#), there were also subglacial lakes under a massively thick ice sheet in Pine Island Bay in the southern Amundsen Sea. The ice there has now dramatically retreated, which allowed us to sample palaeo-lake sediments from aboard Polarstern," says the first author of the *Nature Communications* study, in which the researchers will share their findings today.

"In this regard, we measured the chloride content in the pore water as an indicator of its salinity. In the lower parts of some [sediment cores](#), it was extremely low: a clear sign of freshwater, which originates from lakes

under the ice," explains co-author and AWI geochemist Dr Sabine Kasten. "We modelled the exchange of freshwater in the porous spaces of the sediment with the overlying seawater, to help us reconstruct the dynamics of the transition from a lake to a marine environment," reports Dr José Mogollón from the University of Utrecht. The unique sediments the team collected now provide an archive of changing environmental conditions in the Antarctic and cover a timespan from the present to the Last Glacial Maximum (ca. 21,000 years ago), when the global sea level was roughly 130 metres lower than today.

These lake sediments contain components that are virtually impossible to date, which narrows scientists' ability to estimate their age. "We do know, however, that roughly 11,000 years ago Pine Island Glacier retreated and thinned, causing the moving ice to float on the ocean as an ice shelf. As a result, the lakes that were previously covered by glaciers disappeared into the ocean, when it flooded the Antarctic continental shelf. Only the imprint of the lakes in the form of deep basins and sediments remained preserved, which we have now investigated," says Dr Claus-Dieter Hillenbrand from the British Antarctic Survey.

Satellite-monitoring shows that the movement of water from one [lake](#) to another can cause glaciers draining the Antarctic Ice Sheet to move more quickly. "This aspect needs to be taken into account in models designed to make predictions on the future behaviour and dynamics of ice masses, and with them, the degree to which the sea level will rise," explains AWI marine geologist Kuhn. According to a second study, which Kuhn contributed to and was published in Nature Communications on 17 March 2017, he added: "We have every reason to believe that there are more subglacial lakes in the Antarctic - and more so in the last glacial period - than has been previously assumed. In addition, icecaps like those on the sub-Antarctic island South Georgia and ice sheets reacted much more sensitively and rapidly to climate changes than previously assumed."

**More information:** Kuhn, G., Hillenbrand, C.-D., Kasten, S., Smith, J. A., Nitsche, F. O., Frederichs, T., Wiers, S., Ehrmann, W., Klages, J. P., Mogollón, J. M.: Evidence for a palaeo-subglacial lake on the Antarctic continental shelf. *Nature Communications*. DOI: [10.1038/NCOMMS15591](https://doi.org/10.1038/NCOMMS15591)

Graham, A. G. C., Kuhn, G., Meisel, O., Hillenbrand, C.-D., Hodgson, D. A., Ehrmann, W., Wacker, L., Wintersteller, P., dos Santos Ferreira, C., Römer, M., White, D., Bohrmann, G.: Major advance of South Georgia glaciers during the Antarctic Cold Reversal following extensive sub-Antarctic glaciation. *Nature Communications* 8, 14798, DOI: [10.1038/ncomms14798](https://doi.org/10.1038/ncomms14798)

Provided by Alfred Wegener Institute

Citation: Meltwater lakes under the Antarctic Ice Sheet accelerated glacial retreat in the Earth's past (2017, June 1) retrieved 24 April 2024 from <https://phys.org/news/2017-06-meltwater-lakes-antarctic-ice-sheet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.