


Newly discovered methane consumers in lakes

June 7 2017



Crenothrix bacteria . Credit: Swiss Federal Institute of Aquatic Science and Technology

When decaying organic matter sinks to the bottom of a lake or ocean, methane is produced as the biomass is broken down. Some of the methane is released from the surface into the atmosphere, where it acts as a potent greenhouse gas, while some is broken down by microorganisms in the water column. In a study just published in the

ISME Journal, an international research group reports that, as well as known methane consumers, filamentous bacteria of the genus *Crenothrix* – previously little studied in the environment – are also involved in the methane removal process.

Chance discovery

The role played by *Crenothrix* bacteria was discovered by chance, when the researchers were seeking to quantify [methane](#) removal in the Rotsee (Canton Lucerne) and Lake Zug with the aid of stable isotope labelling. In this method, [methane molecules](#) are labelled with heavy carbon-13 atoms; when ¹³C-labelled methane is assimilated by bacteria, the individual cells can be visualized by means of imaging mass spectrometry. "Typically," says co-author Jana Milucka of the Max Planck Institute in Bremen, "these are small round or rod-shaped cells." In this case, however, the bacteria enriched with ¹³C were not only single round cells but also long, filamentous varieties. This finding was very surprising, because at that point, the researchers didn't know these filamentous bacteria could occur so abundantly in nature. "We then started wondering about their role in the environmental removal of methane," says Milucka.

Eawag geologist Carsten Schubert – an expert on microbial degradation of methane in [water](#) – was also surprised by the group's findings in the two Central Swiss lakes. While the large filamentous *Crenothrix* bacteria have long been known, they are only familiar as a contaminant in drinking water systems, where their proliferation can cause clogging of pipes, sand filters and screens. *Crenothrix* [bacteria](#) have not previously been detected in [lake](#) water, because they have not been specifically sought and they are difficult to identify with molecular genetic methods. Schubert concludes: "We seem to have completely underestimated their role in the biogeochemical cycle."

The researchers have now shown that Crenothrix are not only a stable part of the microbial community in freshwater lakes but may also be the most important methane consumers.

More information: Kirsten Oswald et al. Crenothrix are major methane consumers in stratified lakes, *The ISME Journal* (2017). [DOI: 10.1038/ismej.2017.77](https://doi.org/10.1038/ismej.2017.77)

Provided by Swiss Federal Institute of Aquatic Science and Technology

Citation: Newly discovered methane consumers in lakes (2017, June 7) retrieved 3 May 2024 from <https://phys.org/news/2017-06-newly-methane-consumers-lakes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--