

# Where ocean meets sky—new NASA radar gets a tryout

June 2 2017, by Andrew Good

---



2010 photo of a shoreline in Bay Jimmy, Plaquemines Parish, Louisiana, impacted by the BP Deepwater Horizon oil spill. Oil weakens and kills vegetation, leading to the loss of roots that help hold soil together. Credit: Bruce A. Davis, Department of Homeland Security

Ocean currents and winds form an endless feedback loop: winds blow over the ocean's surface, creating currents there. At the same time, the hot or cold water in these currents influences the wind's speed.

This delicate dance is crucial to understanding Earth's changing climate. Gathering data on this interaction can also help people track oil spills, plan shipping routes and understand ocean productivity in relation to fisheries.

Instruments already exist that measure ocean currents, and others that measure wind, such as NASA's QuickScat and RapidScat. But a new, airborne radar instrument developed by NASA's Jet Propulsion Laboratory in Pasadena, California, is able to measure both.

Called DopplerScat, the instrument is a spinning radar that "pings" the ocean's surface, allowing it to take measurements from multiple directions at once. It's a step up from previous technology, which could simultaneously measure current from one or two directions at the most, and couldn't measure properties of the sea surface as completely as this new instrument.

These measurements would make DopplerScat a valuable addition to future satellite missions, said Ernesto Rodriguez, science lead for the instrument at JPL.

"DopplerScat gives us unprecedented simultaneous observation of wind and currents," Rodriguez said. "Because it combines observations of the surface over a wide swath, we can now take a high-resolution snapshot of the interacting ocean and atmosphere unavailable from previous instruments."


DopplerScat was developed at JPL with funding from NASA's Earth Science Technology Office. As with a highway patrolman's speed gun, it calculates the Doppler effect of a radar signal bouncing off an object. As that object moves closer or farther away, it detects these changes and figures out its speed and trajectory. Those measurements are combined with data from a scatterometer, which detects the reflection of the [radar](#)

[signal](#) from the ocean's surface. The more "scattering" the radar observes, the rougher the waves. From the roughness and orientation of the waves, wind speed and direction can be calculated.

Though it had been tested in two field sites in 2016, DopplerScatt found its ideal proving ground this past April, when the DopplerScatt team joined several agencies conducting scientific research off the U.S. Gulf Coast.

The initiative, called the Submesoscale Processes and Lagrangian Analysis on the Shelf (SPLASH) campaign, was focused on tracking oil spillage and leaks. It was led by the Consortium for Advanced Research on Transport of Hydrocarbon in the Environment (CARTHE), a research team that focuses on how these leaks affect the environment.

SPLASH was designed to look at how oil drifts in the Gulf of Mexico, landing on beaches or affecting water quality at the mouth of the Mississippi River. The CARTHE team's research relied on "drifters"—donut-shaped floats with GPS-units attached.



The DopplerScatt radar at JPL, before being attached to the bottom of a King Air B200 plane. Credit: NASA/JPL-Caltech

The U.S. Naval Research Laboratory, a member of the CARTHE team, provided high-resolution computer modeling to predict the currents and where the drifters would go.

Enter JPL's DopplerScatt team. Rodriguez and Principal Investigator Dragana Perkovic-Martin saw an opportunity to prove the JPL technology's value. Together, the drifters and modeling could provide independent validation of DopplerScatt's measurements, while offering its own unique dataset.

The drifters are limited in that they only collect ocean data, and do so in sparse regions over the course of days. DopplerScatt, fixed to the bottom of a King Air B200 plane, gathered both ocean and wind data over vast areas in just one flyover. It painted a large-scale picture while also validating the Navy's computer models.

"It was basically the first large-scale validation that we've done," Perkovic-Martin said. "The CARTHE team used our data to decide where to place their drifters. In the future, we'll use their data and they'll use ours to improve modeling."

"We were able to study the wind and current in all directions over 16 miles (25 kilometers)," Rodriguez said. "If you scale this up to space, instead of covering Earth once every week, we can cover it once every day."

That kind of accuracy offers more than just real-time tracking of environmental disasters, like [oil spills](#). It could lead to improved forecasts of where that oil will drift and which coastal regions are most at risk. More fundamentally, it could increase our understanding of important mechanisms that govern Earth's weather and climate.

It could also benefit shipping routes, which rely largely on current measurements from buoys.

"The ability to map a coastal region's currents in high resolution would be critical for areas like Alaska, where the currents off a jagged coast are strong and change quickly," Rodriguez said.

Now that the instrument has been validated, Perkovic-Martin said, DopplerScatt is available for use on future NASA airborne science missions.

QuickScat launched in 1999. Despite a partial instrument failure in 2009, it continues to provide calibration data to international scatterometer satellite mission partners. RapidScat ended two years of successful ocean wind monitoring aboard the International Space Station in 2016.

Provided by NASA

Citation: Where ocean meets sky—new NASA radar gets a tryout (2017, June 2) retrieved 25 April 2024 from <https://phys.org/news/2017-06-ocean-skynew-nasa-radar-tryout.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.