


Ruthenium rules for new fuel cells

June 28 2017



Rice University scientists have fabricated a durable catalyst for high-performance fuel cells by attaching single ruthenium atoms to graphene. Credit: Chris Zhang/Rice University

Rice University scientists have fabricated a durable catalyst for high-performance fuel cells by attaching single ruthenium atoms to graphene.

Catalysts that drive the [oxygen reduction reaction](#) that lets fuel cells turn chemical energy into electricity are usually made of platinum, which stands up to the acidic nature of the cell's charge-carrying electrolyte. But platinum is expensive, and scientists have searched for decades for a suitable replacement.

The [ruthenium](#)-graphene combination may fit the bill, said chemist James Tour, whose lab developed the material with his colleagues at Rice and in China. In tests, its performance easily matched that of traditional platinum-based alloys and bested iron and nitrogen-doped graphene, another contender.

A paper on the discovery appears in the American Chemical Society journal *ACS Nano*.

"Ruthenium is often a highly active catalyst when fixed between arrays of four [nitrogen atoms](#), yet it is one-tenth the cost of traditional platinum," Tour said. "And since we are using single atomic sites rather than small particles, there are no buried [atoms](#) that cannot react. All the atoms are available for reaction."

Spreading single ruthenium atoms across a sheet of graphene, the atom-thick form of carbon, turned out to be fairly straightforward, Tour said. It involved dispersing graphene oxide in a solution, loading in a small amount of ruthenium and then freeze-drying the new solution and turning it into a foam.

Baking that at 750 degrees Celsius (1,382 degrees Fahrenheit) in the presence of nitrogen and hydrogen gas reduced the graphene and locked nitrogen atoms to the surface, providing sites where ruthenium atoms could bind.

Materials made at higher and lower temperatures weren't as good, and those made at the proper temperature but without either ruthenium or nitrogen proved the quality of the reaction depended on the presence of both.

The material showed excellent tolerance against methanol crossover and [carbon monoxide poisoning](#) in an acidic medium, both of which degrade

the efficiency of fuel cells; such degradation is a persistent problem with traditional platinum fuel cells.

More information: "Single-Atomic Ruthenium Catalytic Sites on Nitrogen-Doped Graphene for Oxygen Reduction Reaction in Acidic Medium" *ACS Nano* (2017).

pubs.acs.org/doi/abs/10.1021/acsnano.7b02148

Provided by Rice University

Citation: Ruthenium rules for new fuel cells (2017, June 28) retrieved 20 March 2024 from <https://phys.org/news/2017-06-ruthenium-fuel-cells.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--