

# Trash-picking seagulls poop tons of nutrients

June 21 2017

---



Seagulls flock to a landfill to find easy sources of food. A study finds their nutrient-rich droppings may cause major water-quality problems in nearby lakes and reservoirs. Credit: Scott Winton, Duke

At least 1.4 million seagulls feed at landfills across North America, which aside from the nuisance it might pose, is also a threat to the health of nearby waters, a new Duke University study finds.

"We estimate these gulls transport and deposit an extra 240 tons of nitrogen and 39 tons of phosphorus into nearby lakes or reservoirs in North America each year through their feces," said lead author Scott Winton, a 2016 doctoral graduate of Duke's Nicholas School of the Environment.

The added nutrients contained in the birds' droppings can contribute to extensive algal blooms that rob surface waters of much of the oxygen needed to sustain healthy aquatic animal life—a process known as eutrophication.

Oxygen depletion and algal toxins that result from the blooms can have far-reaching ecological and economic impacts, including fish kills, increased costs for local governments, and reduced recreational or fishing values in affected waters.

"It costs local U.S. governments an estimated \$100 million a year in nutrient offset credits to address or prevent the problem and maintain nutrient levels at or below the total maximum daily load threshold for water quality," said Mark River, a doctoral student at Duke's Nicholas School, who conducted the research with Winton.

The scale of the problem and the cumulative cost of dealing with it may be even larger than the new study suggests, said Winton, who is now a visiting postdoctoral fellow at ETHZurich, a science and technology university in Switzerland.

"We estimated and mapped a landfill-gull population of 1.4 million based on documented sightings reported in the eBird Citizen Science database. But the actual population is probably greater than 5 million," Winton said. "That means the amount of nutrients deposited in the lakes, and the costs of preventing or remediating the problem, could be substantially higher."

Winton and River published their study, which is the first to look at the transport of nutrients into [surface waters](#) from seagulls at landfills, on June 15 in the journal *Water Research*.

They conducted the research at landfills near two major drinking water reservoirs—Jordan Lake and Falls Lake - that serve the Raleigh-Durham region of North Carolina. Nitrogen and phosphorus loading data from these two lakes were then scaled up to estimate total loading at water bodies near landfills across North America using a well-established model for measuring the nutrient transport of carnivorous birds.

The findings are applicable to lakes and reservoirs in other parts of the world, as well.

"The idea that gull feces can be a major [water](#) quality problem may sound comical—until you look at data from an individual [lake](#)," Winton said. "In Jordan Lake, for instance, we found that a local flock of 49,000 ring-billed gulls deposit [landfill](#) feces containing nearly 1.2 tons of phosphorus into the lake annually."

That amount, he said, is equivalent to roughly half of the total phosphorus load reduction target established for the New Hope Creek watershed of Jordan Lake. Offsetting this added phosphorus costs local governments about \$2.2 million annually—largely through long-term programs aimed at reducing other sources of inflowing nutrients such as urban stormwater or agricultural runoff.

Reducing the size of the gull flocks may also be an option worth pursuing, Winton said. Reducing the size of open landfills, and covering trash more quickly after it is dumped, are among the ways managers can reduce the gulls' easy access to a food supply and encourage a flock to disperse to other feeding habitats.

"There's a decent history of [local governments](#) implementing gull management at landfills because of fears about airplane strikes or because of the nuisance factor to nearby communities," Winton said. "It might be cost-effective to pursue some of these non-lethal mitigation methods to reduce nutrient loading, as well."

**More information:** R. Scott Winton et al, The biogeochemical implications of massive gull flocks at landfills, *Water Research* (2017). [DOI: 10.1016/j.watres.2017.05.076](https://doi.org/10.1016/j.watres.2017.05.076)

Provided by Duke University

Citation: Trash-picking seagulls poop tons of nutrients (2017, June 21) retrieved 20 April 2024 from <https://phys.org/news/2017-06-trash-picking-seagulls-poop-tons-nutrients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.